

Meeting the Cool New Neighbors V: A 2MASS-Selected Sample of Ultracool Dwarfs

Kelle L. Cruz¹

Department of Physics and Astronomy, University of Pennsylvania, 209 South 33rd Street, Philadelphia, PA 19104; kelle@sas.upenn.edu

I. Neill Reid¹

Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218; and Department of Physics and Astronomy, University of Pennsylvania, 209 South 33rd Street, Philadelphia, PA 19104; inr@stsci.edu

James Liebert¹

Department of Astronomy and Steward Observatory, University of Arizona, Tucson, AZ 85721; liebert@as.arizona.edu

and

J. Davy Kirkpatrick, Patrick J. Lowrance¹

Infrared Processing and Analysis Center, California Institute of Technology, Pasadena, CA 91125; davy@ipac.caltech.edu, lowrance@ipac.caltech.edu

ABSTRACT

We present initial results of our effort to create a statistically robust, volume-limited sample of ultracool dwarfs from the 2MASS Second Incremental Data Release. We are engaged in a multifaceted search for nearby late-type dwarfs and this is the first installment of our search using purely photometric selection. The goal of this work is a determination of the low-mass star and brown dwarf luminosity function in the infrared. Here, we outline the construction of the sample, dubbed 2MU2, and present our first results, including the discovery of 186 M7–L6 dwarfs—47 of these are likely to be within 20 pc of the Sun. These results represent 66% of the ultracool candidates in our sample yet constitute an 127% increase in the number of ultracool dwarfs known within the volume searched (covering 40% of the sky out to 20 pc). In addition, we have identified 10 M4–M6.5 objects that are likely to be within 20 pc (or within 1σ). Finally, based on these initial data, we present a preliminary luminosity function and discuss several

¹Visiting Astronomer, Kitt Peak National Observatory, NOAO, which is operated by AURA under cooperative agreement with the NSF.

interesting features of the partial sample presented here. Once our sample is complete, we will use our measured luminosity function to constrain the mass function of low-mass stars and brown dwarfs.

Subject headings: Galaxy: stellar content—solar neighborhood—stars: low-mass, brown dwarfs—stars: luminosity function, mass function

1. Introduction

Ultracool dwarfs (spectral types M7 and later) include stars and brown dwarfs and can have masses as small as several Jupiter masses. The search for these ultracool dwarfs has been greatly enabled by three recent deep, wide-sky surveys: the Two-Micron All Sky Survey (Skrutskie et al. 1997, 2MASS), the Deep Near Infrared Southern Sky survey (Epchtein et al. 1999, DENIS), and the Sloan Digital Sky Survey (York et al. 2000, SDSS).

Two methods are used to identify late-type dwarfs in these large data sets: direct query of the survey’s photometric data products; and using those data in combination with new and existing proper motion data and optical photometry. The former method has yielded close to 200 L dwarfs and over 180 ultracool M dwarfs in the three surveys (notable for quantity are Phan-Bao et al. (2003, 2001); Schneider et al. (2002); Hawley et al. (2002); Gizis et al. (2000); Kirkpatrick et al. (2000, 1999); Martín et al. (1999)). The other tack has unveiled many previously unknown late-type members of our Solar neighborhood (if not as many L dwarfs). Near-infrared photometry from DENIS and 2MASS has been combined with Luyten’s proper motion catalogs (Luyten (1979, LHS), Luyten (1980, NLTT)), new proper-motion catalogs (Lépine, Shara, & Rich 2002), and with new and existing optical photometry to identify nearly 200 M dwarfs but less than 10 L dwarfs within 35 pc of the Sun (Salim, Lépine, Rich, & Shara 2003; Lépine, Rich, & Shara 2003; Reylé, Robin, Scholz, & Irwin 2002; Lodieu, Scholz, & McCaughrean 2002; Scholz & Meusinger 2002; Reid & Cruz 2002; Reid, Kilkenny, & Cruz 2002; Cruz & Reid 2002; EROS Collaboration et al. 1999).

While both methods have yielded many discoveries, both have biases and limitations that make it difficult to use these data sets to study the new population of ultracool dwarfs statistically. The use of proper motion and optical data has the advantage of finding the nearest objects but this type of study is restricted to bright magnitudes and hence, earlier-type objects. In addition, proper motion searches are strongly biased towards objects with intrinsically high space motions with respect to the Sun. Photometric searches avoid a kinematic bias and probe much larger spatial volumes by reaching fainter magnitudes than proper motion surveys. However, previous photometric searches are characterized by small sky coverage and/or restrictive color-criteria.

Our aim is to use both proper motion and photometric searches to complete the census of the nearest ultracool dwarfs. In the first three papers in this series (Reid & Cruz 2002; Reid, Kilkenny, & Cruz 2002; Cruz & Reid 2002, hereafter Papers I, II, & III) we presented the first results of our

effort to cross-reference the 2MASS Second Incremental Release with the NLTT catalog. While we uncovered nearly 150 new objects within 20 pc, most of them are early and mid-M dwarfs. Reid et al. (2003) (hereafter Paper IV) presented the discovery of an M8.5 dwarf within 6 pc found in the course of the program described in this Paper. Here, we present the initial results and the methodology of our photometric search of the 2MASS Second Incremental Release for nearby objects cooler than spectral type M6.

Building on this foundation, we have created a statistically robust sample of ultracool dwarfs from the 2MASS Second Incremental Release that is complete within 20 pc of the Sun for spectral types M7 to L8. The sample is dubbed 2MU2—2MASS ultracool dwarfs from the Second Data Release. This volume-limited sample permits detailed investigation of the overall range of properties of these low-mass objects which span the stellar/brown dwarf boundary. In particular, once the observations are complete, we will have an infrared luminosity function which we will use to constrain the mass function of low-mass stars and brown dwarfs.

The current knowledge of the field mass function of low-mass stars is based primarily on two projects: the Palomar/Michigan State University Nearby-Star Spectroscopic Survey (Reid, Hawley, & Gizis 1995; Reid, Gizis, & Hawley 2002); and the 8 pc sample (Reid & Gizis 1997; Reid et al. 1999, 2003) (hereafter PMSU and 8 pc sample). Both datasets yield a mass function that is consistent with a power-law distribution ($\Psi(M) = M^{-\alpha}$) with $\alpha = 1.2$ at low masses ($M < 0.6M_{\odot}$) (see Figure 12 of Reid, Gizis, & Hawley (2002)). However, both samples only extend to $0.08M_{\odot}$ —just above the stellar/brown dwarf boundary for solar metallicity objects. Reid et al. (1999) made the first attempt to extend coverage to substellar mass objects in the field, but their study is hampered by sparse statistics. The project described in this Paper is the first concerted effort focused on studying the field mass function in the regime of low-mass stars and brown dwarfs.

This is the fifth paper in a series that report results of our multifaceted study of nearby late-type objects in the immediate Solar neighborhood. While the previous papers in the series have concentrated on objects with large proper motions, here we present our first results of purely photometric selection of candidate nearby dwarfs using the 2MASS survey. In § 2 we describe the creation of the 2MU2 sample. We present our observations in § 3 and the spectral type and absolute magnitude calibrations in § 4. We discuss interesting objects in § 5 and present the characteristics of the portion of the 2MU2 sample presented here and a preliminary luminosity function in § 6. Our conclusions are in § 7.

2. The 2MU2 Sample

The primary goal of this project is to create a sample of ultracool dwarfs within 20 pc using the 2MASS Second Incremental Data Release. This catalog covers 48% of the sky and contains over 162 million point sources with J ($1.25 \mu\text{m}$), H ($1.65 \mu\text{m}$), and K_S ($2.17 \mu\text{m}$) photometry and highly accurate astrometry. While we use several methods to refine the 2MU2 sample, as described in

detail below, there are three primary selection criteria. The first requires targets to have a galactic latitude greater than 10° in order to avoid the Galactic plane. Two additional cuts are based on the JHK_S color-color and color-magnitude sequences of M and L dwarfs with trigonometric parallax measurements. The final 2MU2 sample of viable candidates contains 1225 targets and covers 40% (16,350 sq. deg.) of the sky. Full details of the selection procedures are given in this section.

2.1. Selection Criteria

The initial 2MASS query, using the Gator tool provided by IRSA², required $|b| > 10^\circ$, $J - K_S > 1$, and rejected extended objects. These criteria selected ~ 11.3 million sources. Custom built IDL code was used to further cut the sample in a series of steps which are detailed below and summarized in Table 1.

Cataloged Cloud and Dense Regions: Objects which are associated with star-formation regions (e.g. Orion, Lupus, etc.) were eliminated from the outset. Rough positions and dimensions for those reddening regions were obtained from Dame et al. (1987) and Dutra & Bica (2002). However, since the high density of sources persisted on the fringes of those regions, we enlarged the areas excluded from our sample. The positions of those regions are listed in Table 2. In addition, the dense stellar associations of the LMC, SMC, 47 Tuc, M31, and M33 were also excluded. A total of ~ 1.65 million sources were eliminated based on these, reducing the sample to ~ 9.65 million targets. These cuts are taken into account in computing the total areal coverage of the final 2MU2 sample.

$J/(J - K_S)$: Our goal is to identify nearby objects of spectral types M7 and later. This is accomplished with two cuts in $J/(J - K_S)$. The left panel of Figure 1 is the color-magnitude diagram for the tail end of the main sequence as it would appear if all of the objects were at 20 pc ($M_J + 1.51$). A color-magnitude diagram for a typical 1° region of the 2MASS database is shown in the right panel. We selected objects that meet both of the following criteria:

$$\begin{aligned} (J - K_S) &> 1.0 \\ J &\leq 1.5(J - K_S) + 10.5. \end{aligned} \tag{1}$$

The $(J - K_S) > 1$ criterion eliminates most objects earlier than M7. Objects that are farther away than 20 pc should fall below the line formed by the second criterion while nearby objects lie above it. These criteria have the biggest effect on narrowing our sample by cutting ~ 9.41 million sources, leaving $\sim 236,500$.

One calibrating object (2M 1632, L8) lies just below our line in the JK_S plane and just above the scatter of the main sequence easily visible in the right panel of Figure 1. In our effort to

²<http://irsa.ipac.caltech.edu/>

balance sample completeness with manageable size, we made the JK_S cut such that numerous faint main-sequence stars would be excluded rather than accommodate this blue-ish L8 dwarf.

These criteria were not intended to select for T dwarfs however some early-type examples may fall within our sample (see 2M 0423 in § 5.1).

$(H-K_S)/(J-H)$: To further refine the spectral type distribution of the sample, color cuts were applied in the JHK_S plane. Figure 2 shows the color-color diagram for the same objects as in Figure 1 and our selection criteria:

$$\left. \begin{aligned} (J-H) &\leq 0.8 \\ (J-H) &\leq 1.75(H-K_S) + 0.1875 \\ (J-H) &\geq 1.75(H-K_S) - 0.4750 \end{aligned} \right\} \begin{aligned} &\text{for } 0.30 \leq (H-K) \leq 0.35 \\ &\text{for } 0.35 < (H-K) \leq 1.20. \end{aligned} \quad (2)$$

In addition, the $(J-K_S) > 1$ criterion translates to

$$(J-H) > 1 - (H-K)$$

in JHK_S color-color space. These criteria isolate late-M and L dwarfs and exclude the central region of the giant sequence. This reduced the sample by $\sim 228,000$ objects, leaving 8531 targets.

Figure 2 plots these objects from Kirkpatrick et al. (1999, 2000) whose uncertainties are less than 0.1 magnitudes. About 15% of all of the L dwarfs listed in Kirkpatrick et al. (1999, 2000) scatter outside of our JHK_S cuts. The objects that fall outside of our criteria are among the faintest known L dwarfs with large uncertainties in their colors (between 0.1 and 0.3 magnitudes). Extending the sample to include those regions is not feasible since it would introduce an unmanageable number of unwanted interlopers.

$J/(R-J)$: Included in the 2MASS data products is BR optical photometry for point sources with a counterpart in the Tycho or USNO catalogs within $5''$ of the 2MASS position. For objects where an R magnitude is available, we have applied a cut in $J/(R-J)$ to restrict the sample to objects with optical/NIR colors that are consistent with nearby late-type dwarfs. Figure 3 shows the color-magnitude diagram for cool stars with known parallaxes where the apparent magnitudes have been shifted to as they would appear if at a distance of 20 pc ($M_J + 1.51$). Targets with an available R magnitude ($\sim 40\%$) are required to meet the following criteria:

$$J \leq \begin{cases} 3.50(R-J) + 3.00, & \text{for } (R-J) \leq 1.0, \\ 1.67(R-J) + 4.83, & \text{for } 1.0 < (R-J) \leq 2.8, \\ 5.50(R-J) - 5.90, & \text{for } 2.8 < (R-J) \leq 3.0, \\ 1.16(R-J) + 7.12, & \text{for } 3.0 < (R-J) \leq 5.5. \end{cases} \quad (3)$$

This cut 5337 objects, leaving 3194 targets.

Giants: Bessell & Brett (1988) have shown that some types of giants have colors that meet our JHK_S color criteria and as a result, many M giants persisted in the sample. Figure 2 shows

data for giants in JHK_S color space. Noting that giants tend to have bluer ($H - K_S$) colors and very bright magnitudes ($J < 10$), we are able to get rid of a significant portion of the giants by eliminating objects that meet both of the following criteria. (Note that the previous items have been criteria for inclusion while the following criteria are for exclusion.)

$$\begin{aligned} J &< 10 \\ (J - H) &> 2(H - K_S), \text{ for } 0.375 < (H - K_S) < 0.470. \end{aligned} \tag{4}$$

This cut 698 objects, leaving 2496 targets. The second criterion is shown in Figure 2.

Flags: The 2MASS catalog also provided flags for possible photometric confusion and solar system objects (`cc_flg=000` and `mp_flg=0`). The confusion flag indicates when more than one object is merged and the individual photometry could not be resolved. The solar system flag indicates the point source is associated with a minor planet. Three hundred ten objects were cut based on these flags, leaving 2186.

Uncataloged Reddening Regions: Even though we cut objects at low galactic latitudes and near star-forming regions, there were still several large areas of high density near the plane. These regions are listed in Table 3 and are likely to be reddened sources associated with small/uncataloged molecular clouds, so we have eliminated them. This removed 514, leaving a sample size of 1672 targets.

2.2. Refining the Faint Portion of the 2MU2 Sample

The color-color and color-magnitude diagrams of the 1672 surviving candidates are shown in Figure 4. At this point, the 588 objects with $J \leq 9$ were set aside for separate classification and are discussed in § 2.3. The 1084 objects with $J > 9$ were examined in more detail and 447 objects were eliminated based on the following considerations which are summarized in Table 4:

$(F - J)/(J - K_S)$: Additional optical photometry was obtained from the Guide Star Catalog 2.2 (Morrison, McLean, & GSC-Catalog Construction Team 2001, GSC). $(F - J)$ and $(J - K_S)$ colors were used to eliminate objects that are too blue at optical wavelengths to be ultracool dwarfs. Targets were required to meet the following criteria:

$$(F - J) > \begin{cases} 4.0 \\ 1.67(J - K) + 2.33. \end{cases} \tag{5}$$

This eliminated 137 objects. Recently, Thorstensen & Kirkpatrick (2003), identify one of these eliminated objects as a nearby L dwarf. As suggested by those authors, crowding in the field likely resulted in a mismatch between the 2MASS source and the GSC source and thus an aberrant $(F - J)$ color. All of these 137 objects will be examined by eye to correct for other possible mismatches and will be discussed in a future paper.

Visual Inspection: Two hundred and eleven targets were eliminated via visual inspection of POSS plates and 2MASS images. The images revealed some sources to be artifacts of the 2MASS data, associated with large galaxies or globular clusters, or clearly non-red objects.

SIMBAD: Twenty four objects were eliminated because they are listed in SIMBAD as known carbon stars, pre-main sequence objects, or quasars.

Clouds: Both SIMBAD and visual inspection enabled us to identify 41 objects associated with molecular clouds or reddening regions. These were eliminated.

Bland: Figure 4 shows the color-magnitude and color-color diagrams for the 588 bright objects as well as the surviving faint targets. As can be seen in the left panel there is a high density of faint objects with colors close to $(J - K_S) = 1$. The high density of mid-M dwarfs just bluer than $(J - K_S) = 1$ and the increased photometric uncertainties at fainter magnitudes leads to more scatter into our sample. After many observations of objects in this color space, we were able to confidently eliminate 34 objects with $J - K_S < 1.1$ and $J > 13$ as distant mid-M dwarfs.

These cuts reduced the 2MU2 sample of $J > 9$ objects to 637 ultracool candidates. Table 5 lists the 112 ultracool targets with existing data (predominately from Kirkpatrick et al. (1999, 2000); Gizis et al. (2000)). The remaining 525 require spectroscopic follow-up observations—data for 307 are presented here.

2.3. The Brightest Candidates

Five hundred and eighty-eight of the ultracool dwarf candidates have apparent magnitudes $J \leq 9$. Based on both the color-magnitude distribution, and the fact that most lie relatively close to the Galactic plane, we discuss these sources separately.

Cross-referencing against SIMBAD, using the 2MASS positions, leads to positive identifications for 386 sources, as follows:

- One hundred seventeen sources are within $10\text{--}15''$ of a source from the IRAS catalog. These are likely to be dusty giants, asymptotic giant branch (AGB) stars, supergiants or young protostars. Again, most lie close to the Plane.
- One hundred thirty eight sources lie within $2\text{--}3''$ of stars cataloged in the Henry Draper, Bonner Durchmusterung, Cape Durchmusterung, Cape Photographic Durchmusterung, Guide Star Catalog, or PPM catalog. None of these stars have significant proper motions, and the near-infrared and optical/infrared colors are consistent with red giant or AGB stars.
- Thirteen stars are classified in SIMBAD as carbon stars. Again, all lie on the AGB.

- Eighty stars are identified as Miras and sixteen stars are classed as semi-regular variables; both datasets are predominantly M-type AGB long-period variables. These stars show a strong concentration towards the Plane and the Bulge.
- Eighteen sources are identified with a variety of late-type stars, including red giant variables, symbiotic stars, pre-main sequence variables and a symbiotic star.
- Finally, four sources are matched to known M-dwarf proper motion stars. We discuss these further below.

Near-IR color-magnitude diagrams of these sources are shown in the Appendix.

Two hundred and two bright ultracool candidates have no previous identification listed in the SIMBAD database. However, all save two of those sources have optical counterparts (positional offset of less than $5''$ from the USNOA-1.0 catalog listed in the 2MASS database. Since the USNO catalog is based on scans of the POSS I *O/E* plates in the North (average epoch ~ 1953) and the SERC/UKST *J* and ESO *R* plates in the South (average epoch ~ 1979), this indicates that these sources have proper motions of less than $0''.1 \text{ yr}^{-1}$ and $0''.2 \text{ yr}^{-1}$, respectively—indeed, in every case the measured offsets indicate negligible relative motion (USNO/2MASS). Both sources which lack cataloged USNOA-1.0 counterparts are clearly visible at the 2MASS position on second-epoch UKST IIIaF plates taken in the mid-1980s, also indicating low proper motions.

The presence of non-moving optical counterparts strongly suggests that all of these uncataloged sources are likely to be distant reddened stars or red giants, rather than nearby dwarfs. Confirmation of this hypothesis comes from the $(R - J)/(J - K_S)$ two-color diagram (Figure 5), which shows that the overwhelming majority have significantly bluer $(R - J)$ colors than expected for late-M or L dwarfs. Given the low Galactic latitude of most sources, the potential exists for mismatches. However, as discussed in more detail in the Appendix, spectroscopy of the relatively small number of sources with near-infrared colors consistent with late-M/L dwarfs shows that none are late-type dwarfs.

In summary, only four sources among the 588 ultracool candidates with $J \leq 9$ are likely to be genuine nearby late-type dwarfs. These stars are as follows:

G 180-11: This object is included in the third Catalog of Nearby Stars (Gliese & Jahreiß 1991, pCNS3). Reid, Hawley, & Gizis (1995) measure a spectral type of M4.5 and estimate a distance of $13 \pm 3.9 \text{ pc}$. ($M_V = 13.11$). McCarthy, Zuckerman & Becklin (2001) identify a candidate companion, with $I = 12.6$ and a separation of $1''.5$ at $\text{PA} = 266^\circ$. There is no object at the appropriate location on the POSS I image of this field, suggesting that the fainter object is associated with G 180-11. If the companion were a late-type dwarf, then the absolute magnitude ($M_I \sim 12$) implies a spectral type of $\sim \text{M6}$ and $K_S \sim 10$, ~ 2 magnitudes fainter than the primary. There is no evidence for significant distortion in the 2MASS point-spread function, but that might reflect the large pixel scale ($1''.5$) and consequent poor sampling.

Alternatively, the hypothesized companion might be a white dwarf, in which case the inferred absolute magnitude would be consistent with $T_{eff} \sim 11,000$ K and an expected apparent magnitude of $J \sim 12.5$. Further observations are required to decide between these hypotheses.

G 139-3: This is also a star included in the pCNS3. The spectral type is M4 (Reid, Hawley, & Gizis 1995), and the spectroscopic parallax gives a distance of 13.8 ± 4.0 pc ($M_V = 12.49$).

BD-01 3925D: This object is a wide companion of the K0 dwarf, BD -1:3925A or HD 192263, ($\pi = 50.27 \pm 1.13$ mas; ESA (1997)). BD-01 3925D is not listed in the pCNS3, but has $M_K = 5.61$ for $d = 19.9$ pc, consistent with a spectral type of M0/M1.

EZ Aqr: This object is also known as Gl 866, the well-known triple system (Delfosse et al. 1999). This system is included in pCNS3, has a spectral type of M5.5, and a distance of 3.4 ± 0.03 pc.

With the possible exception of Gl 866ABC, these dwarfs have a relatively unusual location on the $(J - H)/(H - K_S)$ diagram. In the 2MASS All-Sky Release (where the photometry for bright stars is improved over the Second Release), the revised photometry for G 180-11 and G139-3 puts them blueward of our $J - K = 1$ cut but the remaining two objects still have unusually red colors. Since M dwarfs with spectral types earlier than M5 would be expected to have $(J - K_S) < 1.0$, further observations of these objects are desirable. Data for these stars are listed in Table 6.

3. Observations

Follow-up far-red optical spectroscopy has been obtained for 298 potential nearby dwarfs during five separate observing runs at NOAO facilities³. (Data for 9 objects were obtained from other sources.) Tables 7–13 list the positions (as a 2MASS name), photometry, observation date, and derived data for all of the observed targets⁴.

The first run was at the Kitt Peak 2.1 m with the GoldCam spectrograph on 2000 September 29 through October 2. The majority of the run was dedicated to follow-up of proper-motion selected candidates (Paper III). We used a $1''.3$ slit and a 400 line mm^{-1} grating blazed at 8000 Å to give a resolution of 5.1 Å (2.8 pixels) covering the wavelength range 5500 – 9300 Å. Higher orders were blocked using a OG550 filter. Weather was good with $1''$ – $1''.5$ seeing. These data were not flat-fielded due to problems with fringing (see Paper III for a detailed discussion).

Observations were obtained on the Kitt Peak 4 m on 2001 July 13–23 with the RC Spectrograph to cover 6000 – 10000 Å in first order. An OG530 filter was used to block second-order light. We

³Spectra are available upon request from kelle@sas.upenn.edu.

⁴The 2MASS designation is 2MASS Jhhmmss.ss±ddmmss. In addition, we note that the astrometry and photometry for all objects is likely to be different from those listed in the 2MASS All-Sky Release.

achieved a resolution of $5.6 \text{ \AA}$ (2 pixels) with a 316 line mm^{-1} grating blazed at $7500 \text{ \AA}$ and a $1''.5$ slit. The run had fair weather with partly cloudy conditions and a wide range of seeing. We were able to confirm many carbon stars and giants as well as obtain data on brighter targets.

The same instrumental setup was used on 2002 January 21–24 except an OG550 filter was used to block higher orders. Good weather (seeing $0''.9$ – $1''.2$) on the first two nights and the fourth night permitted the use of a $1''$ slit resulting in a resolution of $4.7 \text{ \AA}$ (1.7 pixels). The seeing on the third night ranged from $1''.5$ – $2''$ and a $1''.5$ slit was used to obtain a resolution of $5.2 \text{ \AA}$ (1.9 pixels).

Coincident observing (2002 January 22–25) with the Blanco 4 m on Cerro Tololo also had good weather with mostly $1''$ – $2''$ seeing. A Loral 3K CCD and the RC spectrograph with a 315 line mm^{-1} grating blazed at $7500 \text{ \AA}$ and a $1''$ slit was used to cover the range 5500 – $10000 \text{ \AA}$ with a resolution of $5.5 \text{ \AA}$ (2.8 pixels). An OG515 filter was used to block higher orders.

Immediately following, data were obtained with the CTIO 1.5 m on 2002 January 26–30 with a Loral 1K CCD and RC Spectrograph to cover 6000 – $8600 \text{ \AA}$. Resolution of $6.5 \text{ \AA}$ (3 pixels) was obtained with a 400 line mm^{-1} grating blazed at $8000 \text{ \AA}$ and a $1''.5$ slit. The seeing ranged from $0''.7$ – $2''$.

All the spectra were flat-fielded, corrected for bad pixels, extracted, and wavelength- and flux-calibrated using the standard IRAF packages CCDPROC and DOSLIT. (Data from 2000 September were not flat-fielded.) Wavelength calibration was determined using HeNeAr arcs taken nightly. The flux-calibration was done using HD 19445, HD 84937, Ross 640, G 191-B2b, L 1363-3, and Feige 34 (Oke & Gunn 1983; Hamuy et al. 1994).

4. Results

We have derived spectral types, absolute magnitudes, and distances for all of the observed dwarfs. Absolute magnitudes are estimated based on a spectral type/ M_J relation; and distance estimates are made using the derived M_J and 2MASS J -band photometry. Table 7 lists the data for forty-seven newly discovered objects that lie within 20 pc and have types M7 and later. These objects are additions to the core sample that is used for our luminosity and mass function analysis (see §6. Data for 139 ultracool dwarfs that lie outside of 20 pc are listed in Table 8. Data for ten M dwarfs earlier than M7 with a distance estimates less than 20 pc (or within 1σ) are listed in Table 9 while fifty-three more distant objects are listed in Table 10. Five objects that have spectral features indicative of youth are listed in Table 11, fifty-four giants and carbon stars/dwarfs are discussed below and listed in Tables 12 and 13. Three objects (2MASSI J0150116+152357, 2MASSI J0447575–055324, and 2MASSI J0442007–135623) are reddened, distant, early-type stars. The color-color and color-magnitude diagrams for the 365 dwarfs present in the portion of the 2MU2 sample presented here are shown in Figure 6.

4.1. Spectral Types

We found that using our spectral indices (listed in Table 4 of Paper III) as a predictor of spectral type was unreliable for cool dwarfs. This is especially true for M6–M8 dwarfs since the TiO 5 relation turns around near M7.5 (see Paper III, Figure 3, left panel), leading to highly ambiguous classifications.

As a result, all spectral types for M dwarfs in this Paper are determined via visual comparison with standard star spectra taken during the course of our program. Program objects are typed by being normalized and plotted between spectra from a grid of eight standard M1–M9 dwarfs (Kirkpatrick, Henry, & McCarthy 1991). L dwarf types are determined via comparison with nine standard, integer type L0–L8 publicly available LRIS spectra taken as part of the 2MASS Rare Objects project (Kirkpatrick et al. 1999, 2001). In addition, integer types are favored over half-integer types. The resulting uncertainty for all types is ± 0.5 subtypes, except where noted by a “?” where low signal-to-noise increases the uncertainty to 1 or 2 types.

4.1.1. Giants and Carbon Stars/Dwarfs

In the course of spectroscopic follow-up of potential nearby dwarfs, many targets turned out to be distant giants or carbon stars. In Table 12 we list the photometry, astrometry, and rough spectral types (± 1 type) for the observed giants. The spectral types were estimated by side-by-side comparison with observed spectral standards from Garcia (1989) and spectra kindly provided by J.D. Kirkpatrick from Kirkpatrick, Henry, & Irwin (1997). In Table 13 we list the photometry and astrometry for the carbon stars. Two of these are carbon dwarfs with detectable proper motion between the first epoch sky survey plates and the 2MASS images. These will be discussed in detail in a future paper (P.J. Lowrance et al., in preparation).

4.2. Absolute Magnitudes and Distances

M_J and distance estimates for all of the dwarfs in our sample are listed in Tables 5–10. Absolute magnitudes for spectral types M2–M5.5 were derived using TiO 5, CaH 2, and CaOH as described in Paper III (Figure 4). For later types, however, the index relations are double valued. As a result, we have used spectral type as a predictor of M_J .

Dahn et al. (2002) have shown that M_J is well correlated with spectral type. We have rederived that relation, supplementing data kindly provided by H. Harris with several early objects from PMSU and the 8 pc samples. The data used to make the relation are listed in Table 14 and are plotted in Figure 7. A fourth order polynomial was found to best fit the data, and is valid for types M6–L8:

$$M_J = -4.410 + 5.043(ST) - 0.6193(ST)^2 + 0.03453(ST)^3 - 0.0006892(ST)^4 \quad (6)$$

where ST=0, 10, 18 for spectral types M0, L0, L8, respectively. The uncertainty in spectral type dominates the uncertainty in the estimated M_J .

Distances are derived for program objects using the estimated M_J and 2MASS J photometry. The uncertainties in the derived distances are also dominated by the uncertainty in spectral type. (The uncertainties in the 2MASS J photometry are typically only 0.02–0.03 magnitudes and, as stated above, the uncertainty in M_J is mostly due to the uncertainty in spectral type.)

5. Interesting Individual Objects

Spectra for many of these objects are shown in Figures 8 – 10.

5.1. L Dwarfs within 10 pc

2M 0423: L7.5 at 9.2 pc. This object is typed as T0 by Geballe et al. (2002) based on the strength of water absorption and the detection of methane absorption bands in the 1–2.5 μm region. In addition, Schneider et al. (2002) type this object as L5: based on low-resolution optical spectra. In general, spectral types based on optical and near infrared spectra agree, but there are a few cases where they do not. This issue is discussed in Burgasser, Kirkpatrick, Liebert, & Burrows (2003) and will be addressed in detail in a future paper (J.D. Kirkpatrick et al., in preparation).

2M 0835: L5 at 8.3 pc. This L dwarf is among the brightest of its type, with $J = 13$. As such, it is a good candidate for high resolution observations to check for a possible companion. A trigonometric parallax measurement would also help to confirm this as a single L dwarf within 10 pc or a multiple system at a slightly larger distance.

5.2. Brown Dwarfs

Two very nearby L dwarfs in our sample have Li I absorption at 6708 Å confirming their brown dwarf status.

2M 0652: L4.5 at 11.1 pc with strong lithium absorption feature.

2M 2057: L1.5 at 15.7 pc. This spectrum displays both lithium absorption and $\text{H}\alpha$ emission. Kelu-1 and the previously mentioned 2M 0423 also both have these features (Ruiz, Leggett, & Allard (1997) and J.D. Kirkpatrick et al., in preparation).

5.3. Active Objects

We observed several objects that had unusually strong $H\alpha$ emission—we list the equivalent width (EW) of the line below. The activity level of the entire sample will be discussed in a future paper (J. Liebert et al., in preparation).

LP 423- 31/2M 0752: This M7 (at 10.5 pc) was observed three times during our program: twice in 2002 January and once during a run dedicated to NLTT follow-up in 2001 November. On all occasions, the emission strength was very strong with the EW ranging from 33–45 Å.

2M 1707: Emission in L dwarfs is not common—only two presented here display it. This L0.5 (at 26 pc) has significant $H\alpha$ with EW=35 Å and this object should be monitored to check whether we happened to observe during a period of unusually high activity. A spectrum of this object is shown in Figure 8.

2M 2057: This brown dwarf with $H\alpha$ emission is discussed in § 5.2 above.

2M 2351: This M7 at 35 pc displayed a strong $H\alpha$ emission (EW=51 Å) in 2001 July and, unfortunately, no further observations were taken.

5.4. Young Objects

We have found five objects which have spectral features similar to young (~ 10 Myr) objects found in the TW Hya association (Webb et al. 1999). However, they do not appear to be associated with any known star forming regions. These are listed in Table 11 and shown in Figures 9–10 in between standard dwarf spectra. All show weak CaH absorption (6750–7050 Å) and most show weak K I doublet absorption (7665 and 7699 Å) compared to similar solar age dwarfs. Both of these features are indicative of low gravity and therefore youth.

2M 0253: From the DSS and 2MASS images, we find that this object has moved approximately 6'' over 43 years. This further supports our hypothesis that this is a young nearby dwarf.

2M 0435: In addition to weak CaH absorption, this object displays weak emission in $H\alpha$, K I and Na I. It is about one magnitude brighter than the two TW Hya candidates (Gizis 2002) and is probably within 30 pc. It is in the direction of the nearby star forming region MBM 20; however, a current estimate of the distance to this cloud puts it between 112 and 161 pc—significantly more distant than our estimate for this object (Sandell, Reipurth, & Gahm 1987; Hearty et al. 2000).

2M 0608: This dwarf also shows enhanced VO absorption (7334–7534 and 7851–7973 Å) which is another characteristic of young late-type objects (J.D Kirkpatrick et al., in preparation). It is also fairly near on the sky to 2M 0619 as discussed below.

2M 0619: This object displays the characteristics of a young dwarf. In addition, this object and 2M 0608 are close to each other on the sky and lie fairly close to the plane. However, rough distance estimates of 30 pc for 2M 0608 and 100 pc for 2M 0619 suggest that they are not actually associated.

2M 2234: This dwarf also displays strong $H\alpha$ emission consistent with it being a young, active object.

5.5. Two Blue L Dwarfs

There are two L dwarfs which have unusually blue colors and can easily be seen as outliers in Figure 6. One of these objects, 2MASS J1300425+191235 (L1, 14 pc), was originally discovered by Gizis et al. (2000) and its data are listed in Table 5. The other object, 2MASS J172139+334415 (L3, 15 pc) was discovered as a part of this program and its data are listed in Table 7.

Both of these objects have significant proper motion. Based on POSS plates measurements, we find $(\mu_\alpha, \mu_\delta) = -0.7, -1.1''\text{yr}^{-1}$ for 2M 1300 and $(\mu_\alpha, \mu_\delta) = -1.6, 0.6''\text{yr}^{-1}$ for 2M 1721. Gizis et al. (2000) also pointed out 2M 1300’s unusual colors and high velocity and suggested that it is likely to be old. The large proper motion of 2M 1712 supports their claim, however, we plan to obtain near-infrared spectroscopic observations of these two objects to find out which spectral features are causing the anomalous colors.

5.6. LP 775-31 & LP 655-48

Recently, McCaughrean, Scholz, & Lodieu (2002) pointed out that these two objects (aka 2MASSI 0435161–160657 and 2MASSI 0440232–053008) were mistyped in Paper III and that a retyping places them within 10 pc. While we agree that these objects are cooler than M6 as we previously stated, we find them both to be spectral type M7 (not M7.5 and M8 as in McCaughrean et al.), which puts them at 8.6 and 9.8 pc respectively. Clearly, trigonometric parallaxes are required to determine unambiguous distances to these nearby dwarfs.

6. Sample Characteristics and Preliminary Luminosity Function

In this Paper we present new data on almost 300 objects. These data, combined with the previously known objects, represent 66% of the ultracool candidates in the 2MU2 sample. The color-magnitude and color-color diagrams for those data are shown in Figure 6. In this section, we point out some of the characteristics of the completed portion of the 2MU2 sample that are relevant to the study of nearby low-mass stars and brown dwarfs.

The upper panel of Figure 11 shows the spectral type distribution of the portion of the 2MU2 sample presented here, separately identifying our new discoveries and previously known ultracool dwarfs. We have at least doubled the number of objects in most spectral type bins, and tripled it in others. Not until L5 does the number of previously known objects outnumber our additions. Currently, the follow-up work is the most incomplete for the faintest objects in the sample which is where we expect to find the coolest, faintest objects, which have the latest spectral types.

The distance distribution of the M7–L8 objects presented here is shown in the middle panel of Figure 11. While we have not discovered any objects within five parsecs, we have doubled the number of late-type dwarfs in every other distance bin out to 40 pc. In addition, the number of objects found scales consistently with the increase in volume out to the 20 pc bin. The volume increases 3.4 times from 10 to 15 pc and the number of objects we have found increases 3.2 times. Similarly, from 15–20 pc, the volume increases by a factor of 2.4 and the number of objects included increases by 2.3. This suggests that we are indeed nearly complete out to 20 pc.

In the bottom panel of Figure 11, we show spectral type versus distance. Our current sample includes objects well beyond the 20 pc limit even at spectral types later than L6. This further supports that, once we have complete observations of the full sample including the faintest candidates, we will have identified a complete sample of late-M and L0–L8 dwarfs within 20 pc of the Sun.

Preliminary field luminosity functions in M_J and M_{K_S} (based on the first results from our 2MU2 sample which has a distance limit of 20 pc and an effective sky coverage of 40%) are shown in the top two panels of Figure 12. The components of known binaries are counted separately. M_J is estimated using our spectral type/ M_J relation described in § 4.2 or using 2MASS J band photometry and a trigonometric distance estimate if available. M_{K_S} estimates are obtained by subtracting the 2MASS $J - K_S$ color from M_J .

The faintest bins ($M_J > 14$, $M_{K_S} > 12.5$) should be regarded as substantially incomplete. The number of objects falls off at magnitudes fainter than $M_J = 11$ ($M_{K_S} = 10.5$) as these bins are dominated by evolving brown dwarfs rather than stable stars. While stars remain at a given magnitude for billions of years, brown dwarfs gradually cool to fainter and fainter magnitudes thus depleting the brighter magnitude bins.

The peak at $M_J = 13.75$ with 9 objects is not readily explained. Since we are incomplete at these faint magnitudes ($M_J > 13.5$) and the number of objects in these bins will only go up with increased completeness, it is possible that the peak is not an isolated bump, but rather represents a change to an increasing slope or a plateau of the luminosity function for $M_J > 13.5$. This could be explained as a population of old, massive brown dwarfs that have evolved—essentially, this population has migrated over time from $11.2 < M_J < 13.5$ to $M_J > 13.5$. On the other hand, the peak is not evident in the M_{K_S} luminosity function. It is important to keep in mind that these luminosity functions are based on about two-thirds of our sample of ultracool candidates, which is most lacking in the latest types. While this feature is tantalizing and might have many hypothetical explanations, a luminosity function based on our complete sample, which includes objects down to

$J = 17$, will be better suited to interpretation.

The spectral type distribution of the ultracool dwarfs presented here with distance estimates within 20 pc is shown in the bottom panel of Figure 12. There is a striking lack of L0s—the one object is actually a component of the 2M 0746 system. One possibility for this is that the morphological features corresponding to L0 span a smaller range of temperature than the other types, thus objects evolve through L0 faster and we see fewer objects. However, we do not see a similar lack of L0s in the entire sample (top panel of Figure 11).

The objects presented in this paper represent the brighter part of our 2MU2 sample. While we only present 66% of the ultracool candidates here, of the 219 remaining, we have obtained data on a further 150 objects which will be discussed in a future paper. Forty-eight of the 69 objects without follow-up observations are faint and have proved elusive from 4 m class telescopes. The left panel of Figure 13 shows the color-magnitude distribution of the objects not presented here, while the right panel shows magnitude against spectral type for all of the objects listed in the L Dwarf Archive⁵ along with a rough magnitude limit for our current sample and the 20 pc limit. Based on these, we believe that many nearby late-type L dwarfs are in our 2MU2 sample, but due to their faintness, they still lack follow-up observations. As discussed above, these objects are essential to understanding the luminosity and mass functions of ultracool dwarfs.

Based on initial data reduction of the 150 objects with follow-up observations but not included in this paper, our preliminary luminosity function is $\sim 90\%$ complete for the early-L dwarfs (L0–L4) found within 20 pc. Taking into account our sky coverage of 40%, we find a early-L dwarf space density of 0.0019 pc^{-3} .

7. Conclusions

We have described our project to create a statistically robust, volume-limited survey for ultracool dwarfs (spectral types M7–L8). Our goal is to determine the infrared luminosity function and constrain the mass function of late-type stars and brown dwarfs. We have presented our initial findings—the discovery of 186 new late-M and L dwarfs. Forty-seven of these are additions to the 20 pc nearby star census of ultracool dwarfs, including two confirmed brown dwarfs. These data, combined with previously known nearby objects, are a significant step towards estimating the mass distribution of ultracool field dwarfs. Our future work, especially with the addition of the coolest objects, will illuminate even further the statistical properties of low-mass stars and brown dwarfs in the Solar neighborhood.

We would like to thank the numerous NOAO telescope operators and support staff at Kitt Peak and Cerro Tololo that made this work possible and endured our busy observing program:

⁵<http://spider.ipac.caltech.edu/staff/davy/ARCHIVE/>

Alberto Alvarez, Ed Eastburn, Bill Gillespie, Angel Guerra, Hal Halbedel, Hillary Mathis, and Sergio Pizarro. We would also like to thank Nadya Gorlova and Curtis Cooper for assistance at the 2001 July Kitt Peak observing run. This research was partially supported by a grant from the NASA/NSF NStars initiative, administered by JPL, Pasadena, CA. KLC acknowledges support from a NSF Graduate Research Fellowship. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and IPAC/CalTech, funded by NASA and the NSF; the NASA/IPAC Infrared Science Archive, which is operated by JPL/CalTech, under contract with NASA; and the SIMBAD database, operated at CDS, Strasbourg, France. The Guide Star CatalogII is a joint project of STScI and the Osservatorio Astronomico di Torino. STScI is operated by the AURA, for NASA under contract NAS5-26555. The participation of the Osservatorio Astronomico di Torino is supported by the Italian Council for Research in Astronomy. Additional support is provided by ESO, Space Telescope European Coordinating Facility, the International GEMINI project and the ESA Astrophysics Division.

A. The Brightest Sources

A.1. Introduction

As discussed in the text, 588 sources in our ultracool sample have magnitudes brighter than $J = 9$. Figure 14 shows the distribution of those stars on the celestial sphere and in the near-infrared color-magnitude and color-color planes. Sources with galactic latitudes in the range $-10^\circ < b < 10^\circ$ were excluded *ab initio*. The fiducial dwarf and giant sequences are plotted in the JHK_S two-color diagram, together with data for L dwarfs from Kirkpatrick et al. (1999). Note that we require sources to have $(J - K_S) > 1.0$ and $(H - K_S) > 0.3$; approximately 15% of known L dwarfs have colors outside the limits bounded by the present search criteria.

A.2. Identifications

All 588 sources were cross-referenced against the SIMBAD database using a search radius of 2.0 arcminutes centered on the 2MASS position. As discussed in the text, four stars are confirmed as nearby dwarfs: G 180-11, G 139-3, Gl 866ABC and BD-01 3925D. The results for the remaining 584 candidates are as follows:

IRAS sources: One hundred seventeen 2MASS candidates lie within 10 to 15 arcseconds of a source from the IRAS catalog; given the positional uncertainties of the IRAS astrometry, plus the expectation that IRAS sources should be red in $(J - K_S)$, these are highly likely to be the correct identification for the 2MASS source. The overwhelming majority of these sources are expected to be dusty asymptotic giant branch stars (types M, S and C) or red supergiants. These sources are listed in Table 15. We list J magnitudes for extremely bright

sources where 2MASS H and/or K_S photometry is unavailable, using “:” to denote uncertain measurements. Most of these sources lie close to the Plane, as expected for young giants or luminous AGB stars.

Stellar sources: One hundred thirty eight candidates lie within 2 to 3 arcseconds of stars listed in either the Henry Draper, Bonner Durchmusterung, Cape Durchmusterung, Cape Photographic Durchmusterung, Guide Star Catalog, or PPM catalogues. Data for these sources are given in Table 16. Most of the sample have colors close to the giant sequence in the JHK_S plane, with approximately fifteen stars overlapping with the L dwarf distribution. The latter stars are likely to be carbon stars.

Carbon stars: Thirteen stars in the sample are classed as carbon stars in SIMBAD. Figure 15 shows their distribution on the sky and in the near-infrared plane—there is obvious overlap with the L dwarf distribution in the two-color diagram. Data for these stars are listed in Table 17.

Mira variables: Eighty stars are identified as Miras—M-type long-period variables. Most of these stars lie close to the Galactic Plane, with a particular concentration in the ScoCen region (towards the Bulge). Data are listed in Table 18, and plotted in Figure 15.

Semi-regular variables: Sixteen stars are identified as semi-regular (AGB/RGB) variables. As might be expected, the spatial and color-magnitude distributions are similar to those of the Miras (Figure 15, Table 19), with a strong concentration towards the Plane and the Bulge.

Other: A further eighteen stars are identified as late-type stars based on cross-referencing against SIMBAD. Those stars are listed in Table 20. Most are late-type giants. In particular, StM 218 is from Stephenson’s survey of high-latitude red giants (Stephenson 1986), with additional spectroscopy by Sharples, Whitelock, & Feast (1995); BR B0954-0947 is from the APM QSO survey (Kirkpatrick, Henry, & Irwin 1997); and the DENIS source, from Phan-Bao et al. (2001) is a known red-giant variable. Of the remaining stars, WOH S 11 is listed as type M by SIMBAD, but with no additional information, and TX CVn is a symbiotic binary.

Unmatched sources: Two hundred two sources have no obvious counterpart in the SIMBAD database. 2MASS data for those sources are listed in Table 21, and the spatial and color-magnitude distribution plotted in Figure 16. As discussed in the text, all of these sources have optical counterparts, indicating low proper motions, and most have optical/near-infrared colors which are inconsistent with late-type dwarfs. Given those characteristics, plus the distribution in Galactic coordinates, all are likely to be pre-main-sequence stars or AGB stars.

A.3. Discussion

A.3.1. The Reddest Candidates

The overwhelming majority of the sources listed in Tables 15 to 21 are clearly red giants. Nonetheless, eight sources in Table 16 and twenty-three sources in Table 21 have $(H - K_S) > 0.45$ —colors potentially consistent with L dwarfs. We discuss these sources in detail below:

- The stellar candidates: all have excellent positional agreement between 2MASS and SIMBAD (ICRS) co-ordinates. Three (V355 Gem, CD CVn and BD-08 2741) are known giant stars, and it is extremely likely that the remaining eight are also red giants.

2MASS J0609248+773327: BD+77 225, spectral type K0: $V = 9.49$, so $(V - K_S) = 3.47$, consistent with $M_V \sim 9.0$ and spectral type M0 if it were a dwarf. Those parameters would place it within 10 parsecs of the Sun. However, the low proper motion ($(\mu_\alpha, \mu_\delta) = -12, -0.2 \text{ mas yr}^{-1}$) and the inconsistency between the $(V - K_S)$ color and the observed spectral type indicates that this star is a giant.

2MASS J0700365+260818: GSC 01899-00620, and also identified as V355 Gem and IRAS 06575+2612. This is clearly an evolved star.

2MASS J1300025+472632: CD CVn, listed as $V = 9.39$, $(B - V) = 1.19$, spectral type K0 III and $\pi = 3.31 \pm 1.21$ by SIMBAD. This is a red giant variable, with $(V - K_S) = 3.28$.

2MASS J1341018+563452: HD 238271, $V = 9.55$, $(B - V) = 1.51$, spectral type K5, negligible proper motions ($-7, 6.2 \text{ mas yr}^{-1}$). The low proper motions confirm the star as a red giant, $(V - K_S) = 3.64$.

2MASS J1724215+652915: BD+65 1182, $V = 9.61$, $(B - V) = 1.45$, $(V - K_S) = 3.45$, spectral type K2, negligible proper motions ($-3, -6 \text{ mas yr}^{-1}$). Again, the low motions, measured colors and observed spectral type are most consistent with a giant.

2MASS J1726192+601748: BD+60 1757, $V = 9.94$, $(B - V) = 1.66$, $(V - K_S) = 4.15$, spectral type K2, negligible motions ($-12, 9 \text{ mas yr}^{-1}$). As with 2M 1724, the low motions, measured colors and observed spectral type are most consistent with a giant.

2MASS J1816212+202817: HD 348183, $V = 9.06$, $(B - V) = 1.52$, negligible motions ($-7, -6 \text{ mas yr}^{-1}$), spectral type K7. $(V - K_S) = 3.92$, and another red giant.

2MASS J2015149–153626: GSC 06315-00584, also identified as NSV 12940, a red giant variable.

- The 34 sources with no SIMBAD identification: as noted above, all of these sources have optical counterparts. Genuine ultracool dwarfs are expected to have $(B - R) > 4$ and $(R - K_S) > 7$ for $(H - K_S) > 0.45$ (spectral type later than M8). The corresponding distances (for ultracool dwarfs) are less than 5 parsecs, so the absence of any measured proper motion ($\mu < 0.1 \text{ arcsec yr}^{-1}$) would require transverse velocities of less than 2.5 kms^{-2} .

- 2MASS J0547281–214723:** $(B - R)_{USNO} = 3.0$, $(R - K_S) = 4.2$. Spectroscopy with the CTIO 1.5 m confirms this as a carbon star.
- 2MASS J0613450+522540:** $(B - R)_{USNO} = 2.1$, $(R - K_S) = 6.5$, with no evidence for motion between POSS I and either POSS II or 2MASS. The source appears significantly brighter on the POSS II F plate than on the POSS I E , suggesting variability, and identify this star as a probable AGB variable.
- 2MASS J0620521–164541:** $(B - R)_{USNO} = 1.2$, $(R - K_S) = 7.2$, with no evidence for significant proper motion. CTIO spectroscopy confirms this as an M giant.
- 2MASS J0641403–282102:** $(B - R)_{USNO} = 0.8$, $(R - K_S) = 5.8$, with no evidence for significant proper motion. CTIO spectroscopy confirms this as an M giant.
- 2MASS J0650548–372922:** $(B - R)_{USNO} = 4.1$, $(R - K_S) = 7.9$, with no evidence for significant proper motion. CTIO spectroscopy confirms this as an M giant.
- 2MASS J0652228+452045:** $(B - R)_{USNO} = 1.6$, $(R - K_S) = 5.2$, and no evidence for motion between POSS I and either POSS II or 2MASS. This is likely to be a red giant.
- 2MASS J0657520+662111:** $(B - R)_{USNO} = 1.3$, $(R - K_S) = 5.3$, and no evidence for motion between POSS I and either POSS II or 2MASS. Given the Galactic latitude, $b = +25^\circ$, the colors are most consistent with a reddened background star.
- 2MASS J0658118+263535:** $(B - R)_{USNO} = 5.0$, $(R - K_S) = 4.7$, with no evidence for significant proper motion. CTIO spectroscopy confirms this as an M giant.
- 2MASS J0701322–381421:** $(B - R)_{USNO} = 3.2$, $(R - K_S) = 5.4$, with no evidence for significant proper motion. CTIO spectroscopy confirms this as an M giant.
- 2MASS J0703356–404748:** $(B - R)_{USNO} = 5.9$, $(R - K_S) = 5.5$, $b = -15^\circ$. Highly likely to be a reddened source.
- 2MASS J0710483+305546:** $(B - R)_{USNO} = 4.7$, $(R - K_S) = 4.4$. No motion evident, and clearly fainter on POSS II than POSS I. Likely to be a red giant variable.
- 2MASS J0710574+475818:** $(B - R)_{USNO} = 3.8$, $(R - K_S) = 5.0$, and no evidence for significant proper motion. Likely to be a red giant or reddened background star.
- 2MASS J0721404+194350:** $(B - R)_{USNO} = 1.9$, $(R - K_S) = 4.9$. CTIO spectroscopy identifies this as an M giant.
- 2MASS J0813343–051321:** $(B - R)_{USNO} = 4.0$, $(R - K_S) = 4.9$. CTIO spectroscopy identifies this as a carbon star.
- 2MASS J0829151+182307:** $(B - R)_{USNO} = 4.6$, $(R - K_S) = 4.4$. No evidence for motion; colors strongly suggest highly reddened object.
- 2MASS J1010015–023743:** $(B - R)_{USNO} = 3.1$, $(R - K_S) = 4.4$. CTIO spectroscopy identifies this high-latitude ($b = 41^\circ$) candidate as an M giant.
- 2MASS J1158169–253753:** $(B - R)_{USNO} = 1.6$, $(R - K_S) = 4.3$. CTIO spectroscopy identifies this as another high-latitude ($b = 36^\circ$) M giant.

2MASSI J1502099+593121: $(B - R)_{USNO} = 2.2$, $(R - K_S) = 6.4$, but relatively faint on the POSS II IVN plate ($I \sim 15$) and barely visible on the IIIaJ plate ($B \sim 21$). This is likely to be a high latitude red giant variable.

2MASSI J1502582–355111: $(B - R)_{USNO} = 0.8$, $(R - K_S) = 6.8$. Significantly brighter on the UKST IIIaF plate than on the POSS I 103aE plate, and no evidence for motion. This is likely to be a red giant variable.

2MASSI J1930155–232048: $(B - R)_{USNO} = 0.9$, $(R - K_S) = 8.2$, no evidence for motion and near the Lupus dark cloud. Likely to be a pre-main sequence star or dusty giant.

2MASSI J1942441–295436: $(B - R)_{USNO} = 4.6$, $(R - K_S) = 4.9$, and no evidence for motion. Likely to be a dusty giant.

2MASSI J2025464–163148: $(B - R)_{USNO} = 0.2$, $(R - K_S) = 6.3$, and no evidence for motion POSS I/POSS II/2MASS. Brighter on second epoch UKST IIIaF than IVN, suggesting identification as a red giant variable.

2MASSI J2044540–074359: $(B - R)_{USNO} = 2.1$, $(R - K_S) = 6.8$, and no evidence for motion. Significantly fainter on UKST IIIaF plate, strongly suggesting variability and identification as a red giant.

A.3.2. Cross-checks Against Existing Catalogs

As an additional test, all of the bright ultracool candidates were cross-referenced against the third Catalog of Nearby Stars (Gliese & Jahreiß 1991, pCNS3) and against Luyten’s NLTT proper motion catalog Luyten (1980). The two catalogs were cross-referenced using a search based on position with a search radius of $180''$. There are only twelve matches, including G180-11, G139-3, BD-01 3925D, and EZ Aqr (Gl 866ABC) which are discussed in the main text. In each of the remaining nine cases, it is clear that the 2MASS ultracool source is not the NLTT star.

A.4. Conclusions

The overwhelming majority of the 588 sources in the ultracool sample with $J < 9$ can be eliminated as candidate nearby dwarfs: 386 have previously-cataloged optical or infrared counterparts, and the overwhelming majority of those are AGB stars. Of the remaining 202 sources, only twenty three have colors sufficiently red to be candidate L dwarfs. All of the latter are visible on photographic plate material, and none have either measurable proper motions or optical/near-infrared colors consistent with ultracool dwarfs. We conclude that four proper-motion objects, G 180-11, G139-3, BD-01 3925D, and Gl 866 (Table 6) are the only genuine late-type dwarfs amongst the ultracool candidates with $J \leq 9$.

REFERENCES

- van Altena W.F., Lee J.T., Hoffleit E.D. 1995, General Catalog of Trigonometric Stellar Parallaxes (4th ed.; New Haven:Yale Univ. Obs.)
- Bergeron, P., Leggett, S. K., & Ruiz, M. T. 2001, ApJS, 133, 413
- Bessell, M. S. & Brett, J. M. 1988, PASP, 100, 1134
- Bouy et al., AJ, in press
- Burgasser, A. J., Kirkpatrick, J. D., Liebert, J., & Burrows, A. 2003, ApJ, in press
- Close, L. M., Potter, D., Brandner, W., Lloyd-Hart, M., Liebert, J., Burrows, A., & Siegler, N. 2002a, ApJ, 566, 1095
- Close, L. M., Siegler, N., Freed, M., & Biller, B. 2003, ApJ, 587, 407
- Close, L. M., Siegler, N., Potter, D., Brandner, W., & Liebert, J. 2002b, ApJ, 567, L53
- Cruz, K. L. & Reid, I. N. 2002, AJ, 123, 2828 (Paper III)
- Dahn, C. C. et al. 2002, AJ, 124, 1170
- Dame, T. M. et al. 1987, ApJ, 322, 706
- Delfosse, X., Forveille, T., Udry, S., Beuzit, J.-L., Mayor, M., & Perrier, C. 1999, A&A, 350, L39
- Delfosse, X., Tinney, C. G., Forveille, T., Epchtein, N., Borsenberger, J., Fouqué, P., Kimeswenger, S., & Tiphène, D. 1999, A&AS, 135, 41
- Delfosse, X. et al. 1997, A&A, 327, L25
- Delfosse, X. et al. 2001, A&A, 366, L13
- Dutra, C. M. & Bica, E. 2002, A&A, 383, 631
- Epchtein, N. et al. 1999, A&A, 349, 236 (DENIS)
- EROS Collaboration et al. 1999, A&A, 351, L5
- ESA 1997, The Hipparcos and Tycho Catalogues, ESA SP-1200 (Noordwijk: ESA)
- Fan, X. et al. 2000, AJ, 119, 928
- Freed, M., Close, L. M., & Siegler, N. 2003, ApJ, 584, 453
- Garcia, B. 1989, Bulletin d'Information du Centre de Donnees Stellaires, 36, 27
- Geballe, T. R. et al. 2002, ApJ, 564, 466

- Gizis, J. E. 2002, *ApJ*, 575, 484
- Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., & Burgasser, A. J. 2000, *MNRAS*, 311, 385
- Gizis, J. E., Monet, D. G., Reid, I. N., Kirkpatrick, J. D., Liebert, J., & Williams, R. J. 2000, *AJ*, 120, 1085
- Gizis, J. E., Reid, I. N., Knapp, G. R., Liebert, J., Kirkpatrick, J. D., Koerner, D. W., & Burgasser, A. J. 2003, *AJ*, 125, 3302
- Gliese & Jahreiss, 1991, Third Version of the Catalog of Nearby Stars (Heidelberg: Astron. Rechen-Inst.) (CNS3)
- Hamuy, M., Suntzeff, N. B., Heathcote, S. R., Walker, A. R., Gigoux, P., & Phillips, M. M. 1994, *PASP*, 106, 566
- Hawley, S. L. et al. 2002, *AJ*, 123, 3409
- Hearty, T., Fernández, M., Alcalá, J. M., Covino, E., & Neuhäuser, R. 2000, *A&A*, 357, 681
- Irwin, M., McMahon, R. G., & Reid, N. 1991, *MNRAS*, 252, 61P
- Kendall, T.R., Mauron, N., Azzopardi, M., & Gigoyan, K. 2003, *A&A*, 403, 929
- Kirkpatrick, J. D. et al. 1999, *ApJ*, 519, 802
- Kirkpatrick, J. D. et al. 2000, *AJ*, 120, 447
- Kirkpatrick, J. D., Henry, T. J., & Irwin, M. J. 1997, *AJ*, 113, 1421
- Kirkpatrick, J. D., Henry, T. J., & McCarthy, D. W. 1991, *ApJS*, 77, 417
- Kirkpatrick, J. D., Liebert, J., Cruz, K. L., Gizis, J. E., & Reid, I. N. 2001, *PASP*, 113, 814
- Kirkpatrick, J. D., McGraw, J. T., Hess, T. R., Liebert, J., & McCarthy, D. W. 1994, *ApJS*, 94, 749
- Koerner, D. W., Kirkpatrick, J. D., McElwain, M. W., & Bonaventura, N. R. 1999, *ApJ*, 526, L25
- Liebert, J., Kirkpatrick, J. D., Cruz, K. L., Reid, I. N., Burgasser, A., Tinney, C. G., & Gizis, J. E. 2003, *AJ*, 125, 343
- Lépine, S., Rich, R. M., & Shara, M. M. 2003, *AJ*, 125, 1598
- Lépine, S., Shara, M. M., & Rich, R. M. 2002, *AJ*, 124, 1190
- Lodieu, N., Scholz, R.-D., & McCaughrean, M. J. 2002, *A&A*, 389, L20

- Luyten, W. J. 1979, Catalog of Stars with Proper Motions exceeding $0''.5$ Annually (Minneapolis: University of Minnesota Press) (LHS)
- Luyten, W. J. 1980, Catalog of Stars with Proper Motions Larger than $0''.2$ (Minneapolis: University of Minnesota Press) (NLTT)
- Luyten, W. J. & Ebbighausen, E. G. 1936, AJ, 45, 188
- Martín, E. L., Brandner, W., & Basri, G. 1999, Science, 283, 1718
- Martín, E. L., Delfosse, X., Basri, G., Goldman, B., Forveille, T., & Zapatero Osorio, M. R. 1999, AJ, 118, 2466
- McCarthy, M. F., Bertiau, F. C., & Treanor, P. J. 1964, Ricerche Astronomiche, 6, 571
- McCarthy, Zuckerman & Becklin, AJ, 121, 3259
- McCaughrean, M. J., Scholz, R.-D., & Lodieu, N. 2002, A&A, 390, L27
- Monet, D. G., Dahn, C. C., Vrba, F. J., Harris, H. C., Pier, J. R., Luginbuhl, C. B., & Ables, H. D. 1992, AJ, 103, 638
- Morrison, J. E., McLean, B., & GSC-Catalog Construction Team, I. 2001, AAS/Division of Dynamical Astronomy Meeting, 32
- Oke, J. B. & Gunn, J. E. 1983, ApJ, 266, 713
- Phan-Bao, N. et al. 2001, A&A, 380, 590
- Phan-Bao, N. et al. 2003, A&A, 401, 959
- Reid, I. N. & Cruz, K. L. 2002, AJ, 123, 2806 (Paper I)
- Reid, L. N. & Gilmore, G. 1981, MNRAS, 196, 15P
- Reid, I. N. & Gizis, J. E. 1997, AJ, 113, 2246
- Reid, I. N., Gizis, J. E., & Hawley, S. L. 2002, AJ, 124, 2721
- Reid, I. N., Gizis, J. E., Kirkpatrick, J. D., & Koerner, D. W. 2001, AJ, 121, 489
- Reid, I. N., Hawley, S. L., & Gizis, J. E. 1995, AJ, 110, 1838 (PMSU)
- Reid, I. N., Kilkenny, D., & Cruz, K. L. 2002, AJ, 123, 2822 (Paper II)
- Reid, I. N. et al. 1999, ApJ, 521, 613
- Reid, I. N. et al. 2003, AJ, 125, 354 (Paper IV)
- Reylé, C., Robin, A. C., Scholz, R.-D., & Irwin, M. 2002, A&A, 390, 491

- Ruiz, M. T., Leggett, S. K., & Allard, F. 1997, *ApJ*, 491, L107
- Ruiz, M. T., Wischnjewsky, M., Rojo, P. M., & Gonzalez, L. E. 2001, *ApJS*, 133, 119
- Salim, S., Lépine, S., Rich, R. M., & Shara, M. M. 2003, *ApJ*, 586, L149
- Sandell, G., Reipurth, B., & Gahm, G. 1987, *A&A*, 181, 283
- Schneider, D. P. et al. 2002, *AJ*, 123, 458
- Scholz, R.-D. & Meusinger, H. 2002, *MNRAS*, 336, L49
- Sharples, R. M., Whitelock, P. A., & Feast, M. W. 1995, *MNRAS*, 272, 139
- Sandage, A. & Kowal, C. 1986, *AJ*, 91, 1140
- Skrutskie, M. F. et al. 1997, in *The Impact of Large Scale Near-IR Sky Surveys*, ed. F. Garzon et al. (Dordrecht: Kluwer Academic Publishing Co.), 25 (2MASS)
- Stephenson, C. B. 1986, *ApJ*, 301, 927
- Thorstensen, J. R. & Kirkpatrick, J. D. 2003, *PASP*, in press.
- Tinney, C. G. 1996, *MNRAS*, 281, 644
- Tinney, C. G., Mould, J. R., & Reid, I. N. 1993, *AJ*, 105, 1045
- Tinney, C. G., Reid, I. N., Gizis, J., & Mould, J. R. 1995, *AJ*, 110, 3014
- Webb, R. A., Zuckerman, B., Platais, I., Patience, J., White, R. J., Schwartz, M. J., & McCarthy, C. 1999, *ApJ*, 512, L63
- Wilson, J. 2002, Ph.D. thesis, Cornell University.
- York, D. G. et al. 2000, *AJ*, 120, 1579 (SDSS)




Fig. 1.— Color-magnitude diagram for low-mass stars with trigonometric parallax measurements shifted to 20 pc (*left panel*) and a typical 1° 2MASS field (*right panel*) with our selection criteria. Triangles are from the 8 pc sample. Data for ultracool dwarfs (M7–L8, filled circles) and T dwarfs (filled five-point stars) are from Dahn et al. (2002). Targets are selected if they lie above and to the right of the cuts.




Fig. 2.— Color-color diagram for the same data (plotted with the same symbols) as Figure 1 and our selection criteria. In addition to objects with trigonometric parallax data, we show L dwarfs from Kirkpatrick et al. (1999, 2000) with uncertainties less than 0.1 magnitudes as crosses and open squares are giants. We also show the dwarf and giant sequences (both from Bessell & Brett (1988), transformed to the 2MASS system). Targets are selected if they lie within the enclosed region. The region where bright objects ($J < 10$) are eliminated as giants is enclosed with a dotted line.




Fig. 3.— Color-magnitude diagram for GKM dwarfs (crosses) and L dwarfs (solid circles) with known parallaxes and shifted to 20 pc. The solid line shows the cuts. Targets are selected if they lie above the cut.




Fig. 4.— Color-magnitude and color-color diagrams for the targets that survived the cuts described in § 2.1. Objects fainter than $J = 9$ are candidates for spectroscopic follow-up. The population of objects with $J \leq 9$ are discussed in § 2.3.




Fig. 5.— The $(R - J)/(J - K_S)$ distribution for main-sequence FGKM stars (crosses), L dwarfs (open circles) and bright ultracool candidates with no counterpart listed by SIMBAD (solid triangles).




Fig. 6.— Color-magnitude and color-color diagrams for the all of the cool dwarfs present in this Paper. Crosses are early and mid-M dwarfs (M0–M6.5), filled triangles are late-type M dwarfs (M7–M9.5), and filled circles are L dwarfs. The two blue L dwarf outliers are discussed in § 5.5




Fig. 7.— M_J /spectral type calibration. The solid line is our fourth order fit to the data, while the dashed line is the linear fit found by Dahn et al. (2002) to a similar dataset. Data plotted are listed in Table 14




Fig. 8.— Spectra for some of the interesting objects. Each object is discussed in § 5. The bottom spectrum is not offset and the zero point for each offset spectrum is shown by a dashed line.




Fig. 9.— Spectra for candidate young objects and the M6 and M7 spectral standards LHS 1326 and VB 8. The spectral features of each object are discussed in §5.4. The bottom spectrum is not offset and the zero point for each offset spectrum is shown by a dashed line.




Fig. 10.— Spectra of 2M 0608 and the M9 standard LHS 2065. This object is discussed in §5.4. The bottom spectrum is not offset and the zero point of the offset spectrum is shown by a dashed line.




Fig. 11.— Stacked histogram of the spectral type of all the dwarfs M5 and later in our sample (*top panel*) and distance distributions of the M7–L8 dwarfs (*middle panel*). Darkly shaded region indicates objects previously known while the lightly shaded region represents additions. *Bottom panel* shows spectral type versus distance for dwarfs later than M5. Solid circles are new objects while open triangles are previously known objects, and the solid horizontal line marks our distance limit. Multiplicity has been ignored in all three plots.




Fig. 12.— Preliminary field luminosity function (*top panel*), and spectral type distribution (*bottom panel*) for dwarfs within 20 pc. Hatched region indicates objects with trigonometric parallaxes while shading indicates M_J is based on spectral type. M_J bins are 0.5 mags wide and are labeled with the centroid. Spectral type bins contain both the integer type and the 0.5 class cooler subtype. Multiplicity has been taken into account and the magnitudes corrected.




Fig. 13.— Color-magnitude for the all of the targets in the sample that are not presented here (*left panel*) and J versus spectral type for all 250 objects listed in the L Dwarf Archive (*right panel*). Crosses are objects for which we have data while filled circles represent objects that still require follow-up observations. The dashed line shows where our current incompleteness becomes significant and the solid line marks the 20 pc limit.




Fig. 14.— The (α, δ) and near-infrared $J/(J - K_S)$ and $(J - H)/(H - K_S)$ distributions for all 588 sources in the 2MASS bright ultracool sample. Zero hours of RA is on the left and 12 hours is in the center. The dotted lines in the JHK_S plane mark the selection criteria; the dashed line plots the fiducial giant sequence; the solid line marks the dwarf sequence; and the open circles plot data for L dwarfs from Kirkpatrick et al. (1999).




Fig. 15.— The (α, δ) and near-infrared $J/(J - K_S)$ and $(J - H)/(H - K_S)$ distributions for 2MASS ultracool sources matched against known carbon stars (crosses, Table 17), Miras (open squares, Table 18), semi-regular variables (solid points, Table 19) and other late-type stars (open triangles, Table 20). The lines of RA and the lines in the JHK_S plane are the same as the previous figure.




Fig. 16.— The (α, δ) and near-infrared $J/(J-K_S)$ and $(J-H)/(H-K_S)$ distributions for ultracool sources with no counterpart listed by SIMBAD. The lines of RA and the lines in the JHK_S plane are the same as the previous figure.

Table 1. Accounting of Steps to Create the 2MU2 Sample

Item	Number
2MASS Hits	11,316,668
Clouds	1,271,555
LMC, SMC, & 47 Tuc	383,472
M31 & M33	10,292
J , $(J - K)$	9,414,914
$(J - H)$, $(H - K)$	227,904
R , $(R - J)$	5337
Giants	698
Flags	310
Reddening Regions	514
Total	1672

Note. — The sample started as ≈ 11 million 2MASS hits. The following (non-bold face) entries list reasons for eliminating objects from the sample and their contribution to reducing the final sample size to 1672 candidates. An explanation of each reason for elimination is given in the text.

Table 2. Cataloged Clouds

Region	l_{min}	l_{max}	b_{min}	b_{max}
Cepheus	100	120	11	22
Per OB2	154	163	−25	−10
Taurus	163	202.5	−22	−10
Orion B	202.5	210	−21	−10
Orion A	210	218	−21	−14
Mon R2	210	218	−14	−10
Cham	295	305	−20	−12
Lupus	333	350	10	22
ρ Oph	350	2	13	24
	350	13	10	19
Upper Sco	355	363	16	27
R CrA	4	15	−15	−10
	352	4	−22	−20

Note. — These regions have been eliminated from the sample. The positions have been expanded upon from regions listed by Dame et al. (1987) and Dutra & Bica (2002). Some of the regions overlap.

Table 3. Uncataloged Reddening Regions

l_{min}	l_{max}	b_{min}	b_{max}
0	96	−16	16
150	180	10	13
180	360	−13	13
199	214	−13	−27
308	310	13	16

Note. — Objects lying in these regions have been removed from the sample.

Table 4. Accounting of Steps to Refine $J > 9$ Portion of the 2MU2 Sample

Item	Number
Faint Candidates	1084
$(F - J)/(J - K)$	137
Visual Inspection	211
SIMBAD	24
Clouds	41
Bland	34
Ultracool Candidates	637

Note. — Similar to Table 1 except listing the cuts applied only to the fainter ($J > 9$) portion of the sample and their contribution to reduce the number of ultracool candidates to 637.

Table 5. Previously Known Cool Dwarfs Recovered in the 2MU2 Sample

2MASS Designation	Other Names	J	$J - H$	$J - K_S$	Spectral Type	M_J	d (pc)	$M_{J\pi}$	d_π (pc)	π Ref.	Mult. Ref.	Discovery Ref.
0024246–015819*	BRI 0021–0214	11.860 ^a	0.740	1.280	M9.5	11.6±0.1	11.3±0.7	11.56±0.10	11.5±0.5	1		1
0024442–270825	LHS 1070/LP 881- 64	9.262	0.730	1.030	M5.5	...	...	9.92±0.20	7.39±0.66	2		2
0027559+221932*	LP 349- 25	10.608	0.638	1.047	M8	11.2±0.2	7.80±0.6					3
0030300–145033		16.792	1.437	2.414	L7	14.4±0.2	29.4±3.4					4
0051107–154417		15.230	1.082	1.806	L3.5	12.9±0.2	29.5±3.0					4
0052546–270559	RG 0050.5–2722	13.611	0.617	1.072	M8	11.2±0.2	31.0±2.6	11.94±0.47	21.6±4.7	3		5
0058425–065123		14.320	0.875	1.413	L0	11.7±0.1	33.0±2.1					4
0103320+193536		16.264	1.383	2.111	L6	14.0±0.2	28.1±3.2					4
0104376+145724		13.696	0.672	1.088	M8	11.2±0.2	32.2±2.7					6
0105190+140740		13.588	0.665	1.040	M7	10.7±0.3	37.2±4.4					6
0127391+280553	CTI 012657.5+280202	14.018	0.680	1.188	M8.5	11.3±0.2	34.6±2.5	11.44±0.05	32.8±0.5	4		7
0145452+130600	LHS 1294/LP 468-199	12.783	0.695	1.028	M1.5	...	...					3
0149089+295613		13.443	0.852	1.434	M9.5	11.6±0.1	23.4±1.5	11.68±0.05	22.5±0.4	4		8
0205034+125142		15.680	1.234	2.022	L5	13.6±0.2	26.6±3.0					4
0205293–115930**	DENIS-P J0205.4–1159	14.581	0.991	1.599	L7	(14.4±0.2)	(10.6±1.0)	13.10±0.07 ^b	19.8±0.6	4	1,2	9
0208183+254253		14.015	0.905	1.436	L1	12.0±0.1	25.3±1.7					4
0208236+273740		15.701	1.154	1.833	L5	13.6±0.2	26.9±3.1					4
0220181+241804		13.008	0.686	1.096	M6	10.1±0.4	37.9±6.5					6
0240295+283257		12.679	0.701	1.103	M7.5	11.0±0.2	22.0±2.2					6
0248410–165121*	LP 771- 21/BR 0246–1703	12.557	0.701	1.148	M8	11.2±0.2	19.1±1.6	11.51±0.19	16.2±1.4	3		3
0253202+271333*		12.504	0.667	1.049	M8	11.2±0.2	18.6±1.5					6
0255035–470050*	DENIS-P J0255–4700	13.225	1.036	1.698	L8	14.8±0.1	4.90±0.3					10
0320596+185423*	LP 412- 31	11.744	0.701	1.172	M8	11.2±0.2	13.1±1.1	10.94±0.04	14.5±0.1	4		3
0337036–175807		15.594	1.182	2.006	L4.5	13.3±0.2	28.5±3.2					4
0339352–352544*	LP 944- 20/BRI 0337–3535	10.748	0.731	1.223	M9	11.5±0.1	7.20±0.5	12.25±0.05	5.00±0.1	3		3
0350573+181806	LP 413- 53	12.951	0.729	1.188	M8 ^c	11.2±0.2	22.9±1.9					3
0351000–005244*	LHS 1604/LP 593- 68	11.262	0.670	1.071	M7.5 ^d	11.0±0.2	11.5±1.1	10.43±0.43	14.7±2.9	5		3
0354013+231633		13.122	0.704	1.147	M8.5	11.3±0.2	22.9±1.7					6
0409095+210439		15.545	1.084	1.703	L3	12.7±0.2	37.5±3.6					4
0740096+321203		16.167	1.330	1.984	L4.5	13.3±0.2	37.1±4.3					4
0741068+173845*	LHS 1937/LP 423- 14	11.995	0.633	1.026	M7	10.7±0.3	17.9±2.1					3
0746425+200032**		11.742	0.743	1.255	L0.5	(11.9±0.1)	(9.50±0.6)	11.31±0.03 ^e	12.21±0.04	4	3,2	4
0810586+142039*		12.714	0.669	1.104	M9	11.5±0.1	17.8±1.2					6
0818580+233352*		12.137	0.635	1.007	M7	10.7±0.3	19.1±2.2					6
0820299+450031		16.294	1.295	2.060	L5	13.6±0.2	35.3±4.2					4
0825196+211552*		15.116	1.328	2.071	L7.5	14.6±0.2	12.5±1.0	14.97±0.05	10.7±0.1	4		4

Table 5—Continued

2MASSI Designation	Other Names	J	$J - H$	$J - K_S$	Spectral Type	M_J	d (pc)	$M_{J\pi}$	d_π (pc)	π Ref.	Mult. Ref.	Discovery Ref.
0829066+145622		14.716	0.926	1.599	L2	12.3±0.2	30.3±2.4					4
0832045−012835		14.127	0.818	1.440	L1.5	12.1±0.2	24.9±1.8					4
0840297+182409	LHS 2034/LP 425-140	11.052	0.647	1.003	M6	10.1±0.4	15.4±2.6	10.32±0.04	14.0±0.2	6		3
0850359+105715		16.460	1.234	1.999	L6	(14.0±0.2)	(30.7±3.7)	14.42±0.23	25.6±2.3	4	3,2	8
0853362−032932*	LHS 2065/LP 666- 9	11.185	0.717	1.213	M9	11.5±0.1	8.80±0.6	11.54±0.04	8.50±0.1	6		3
0909575−065818	DENIS-P J0909−0658	13.897	0.810	1.350	L0	11.7±0.1	27.1±1.7					10
0928397−160312		15.337	1.036	1.700	L2	12.3±0.2	40.3±3.2					4
0952219−192431*		11.877	0.595	1.031	M7	10.7±0.3	16.9±2.0					6
1016347+275149*	LHS 2243/LP 315- 53	11.951	0.657	1.005	M7.5	11.0±0.2	15.8±1.5					3
1024099+181553*		12.242	0.664	1.037	M7	10.7±0.3	20.0±2.4					6
1029216+162652		14.307	0.961	1.701	L2.5	12.5±0.2	23.2±2.0					4
1035245+250745		14.700	0.819	1.420	L1	12.0±0.1	34.7±2.4					4
1047126+402643	LP 213- 67	11.417	0.640	1.017	M8	(11.2±0.2) ^f	(11.3±0.9) ^f				4,2	3
1047310−181557	DENIS-P J1047−1815	14.196	0.771	1.299	L2.5:	12.5±0.2	22.0±1.9					10
1049414+253852		12.398	0.648	1.004	M6	10.1±0.4	28.6±4.9					6
1058478−154817*	DENIS-P J1058−1548	14.184	0.944	1.673	L3	12.7±0.2	20.0±1.9	12.99±0.05	17.3±0.3	4		9
1108307+683017*		13.139	0.912	1.539	L1	12.0±0.1	16.9±1.1					6
1112256+354813	Gl 417B	14.573	1.100	1.879	L4.5	(13.3±0.2)	(17.8±1.9)	12.89±0.06	21.7±0.4	7	5,6,7	4
1121492−131308**	LHS 2397a/LP 732- 94	11.929	0.672	1.206	M8.5	(11.3±0.2)	(13.2±1.0)	11.15±0.07 ^g	14.3±0.4	2	8	3
1127534+741107		13.059	0.692	1.088	M8	(11.2±0.2)	(24.0±2.0)				2,7	6
1146344+223052		14.166	0.967	1.525	L3	(12.7±0.2)	(19.9±1.8)	11.99±0.06	27.2±0.6	4	1,3,2	8
1200329+204851		12.850	0.605	1.026	M7	10.7±0.3	26.5±3.1					6
1224522−123835*	BR 1222−1221	12.564	0.733	1.193	M9	11.5±0.1	16.6±1.1	11.40±0.14	17.1±1.1	3		11
1228152−154734	DENIS-P J1228−1547	14.375	1.012	1.570	L5	(13.5±0.2)	(14.6±1.6)	12.85±0.09	20.2±0.8	4	9,1,2	9
1237270−211748		12.665	0.616	1.020	M6	10.1±0.4	32.3±5.6					6
1239272+551537		14.670	1.131	1.927	L5	(13.6±0.2) ^h	(16.7±1.8) ^h				7,6	4
1246467+402715		15.002	1.025	1.699	L4	13.1±0.2	24.1±2.5					4
1246517+314811	LHS 2632/LP 321-222	12.255	0.667	1.022	M6.5	10.5±0.3	22.9±3.3					3
1253124+403403*	LHS 2645/LP 218- 8	12.177	0.620	1.004	M7.5	11.0±0.2	17.5±1.7					3
1300425+191235*		12.710	0.641	1.105	L1	12.0±0.1	13.9±0.9					6
1305401−254106*	Kelu-1	13.417	1.030	1.691	L2	12.3±0.2	16.7±1.3	12.06±0.09	18.7±0.7	4		12
1311391+803222		12.814	0.671	1.100	M8	(11.2±0.2)	(21.5±1.8)				2,7	6
1338261+414034		14.217	0.913	1.466	L2.5	12.5±0.2	22.2±1.9					4
1343167+394508		16.177	1.325	2.066	L5	13.6±0.2	33.5±3.8					4
1403223+300754*		12.691	0.683	1.065	M8.5	11.3±0.2	18.8±1.4					6
1411175+393636		14.679	0.867	1.405	L1.5	12.1±0.2	32.1±2.4					4

Table 5—Continued

2MASS Designation	Other Names	J	$J - H$	$J - K_S$	Spectral Type	M_J	d (pc)	$M_{J\pi}$	d_π (pc)	π Ref.	Mult. Ref.	Discovery Ref.
1426316+155701		12.868	0.686	1.159	M9	(11.5±0.1) ⁱ	(19.1±1.3) ⁱ				10,11,2,7	6
1430378+594325	LHS 2930/LP 98- 79	10.765	0.646	1.009	M6.5	10.5±0.3	11.5±1.6	10.85±0.04	9.60±0.1	6		3
1448033+155414	LHS 2980/LP 441- 17	12.481	0.629	1.010	M6.5	10.5±0.3	25.4±3.5					3
1456383−280947*	LHS 3003/LP 914- 54	9.957	0.630	1.040	M7	10.7±0.3	7.00±0.8	10.96±0.08	6.30±0.2	3		3
1457396+451716		13.145	0.733	1.222	M9	11.5±0.1	21.7±1.4					6
1500342−005944	TVLM 868-54745	12.639	0.656	1.018		10.2±0.7 ^j	31.3±9.8 ^j					11
1506544+132106*		13.414	1.002	1.666	L3	12.7±0.2	14.1±1.3					6
1507476−162738*		12.822	0.920	1.520	L5	13.6±0.2	7.10±0.8	13.50±0.03	7.33±0.03	4		4
1543581+320641	LP 328- 36	12.731	0.615	1.004	M6.5	10.5±0.3	28.5±4.0					3
1546054+374946*		12.437	0.644	1.018	M7.5	11.0±0.2	19.7±1.9					6
1551066+645704		12.870	0.721	1.135	M8.5	11.3±0.2	20.4±1.5					6
1553199+140033		13.019	0.750	1.172	M9	11.5±0.1	20.5±1.3					6
1615441+355900		14.571	1.020	1.630	L3	12.7±0.2	24.0±2.2					4
1627279+810507		13.042	0.710	1.168	M9	11.5±0.1	20.7±1.4					6
1635191+422305		12.886	0.676	1.086	M8	11.2±0.2	22.2±1.8					6
1658037+702701*		13.309	0.766	1.390	L1	12.0±0.1	18.3±1.2	11.96±0.04	18.6±0.2	4		6
1726000+153819		15.653	1.193	2.017	L2	12.3±0.2	46.6±3.9					4
1728114+394859		15.964	1.183	2.066	L7	(14.4±0.2) ^k	(20.1±1.9) ^k				7,6	4
1733189+463359		13.214	0.806	1.357	M9.5	11.6±0.1	21.0±1.3					6
1750129+442404		12.791	0.622	1.031	M7.5	(11.0±0.2)	(23.2±2.3)				2	6
1757154+704201*	LP 44-162	11.446	0.604	1.073	M7.5	11.0±0.2	12.5±1.2					3
1843221+404021*	LHS 3406/LP 229- 30	11.299	0.632	1.030	M8 ^l	11.2±0.2	10.7±0.9	10.55±0.04	14.1±0.2	6		3
2049197−194432		12.871	0.635	1.106	M7.5	11.0±0.2	24.1±2.3					6
2113029−100940		12.863	0.642	1.049	M6	10.1±0.4	35.4±6.1					6
2140293+162518		12.943	0.673	1.164	M8.5	(11.3±0.2)	(21.1±1.5)				11,2,7	6
2147436+143131		13.842	0.708	1.190	M8	(11.2±0.2)	(34.5±2.9)				7,6	6
2202112−110945	LP 759- 17/DENIS-P J2202−1109	12.364	0.650	1.007	M6.5	10.5±0.3	24.1±3.4					3
2206228−204705		12.381	0.677	1.056	M8	(11.2±0.2) ^m	(17.6±1.5) ^m				11,2,7	6
2206449−421720		15.569	1.091	1.974	L2	12.3±0.2	44.9±3.8					4
2224438−015852*		14.052	1.249	2.035	L4.5	13.3±0.2	14.0±1.5	13.77±0.04	11.4±0.1	4		4
2234139+235955		13.176	0.823	1.341	M9.5	11.6±0.1	20.7±1.3					6
2235490+184029	LP 460- 44	12.458	0.632	1.125	M7	10.7±0.3	22.1±2.6					3
2255584+282246	LP 345- 18	12.554	0.619	1.016	M6	10.1±0.4	30.7±5.3					3
2306292−050227*		11.372	0.654	1.084	M7.5	11.0±0.2	12.1±1.2					6
2331016−040619		12.937	0.648	1.007	M8	(11.2±0.2)	(22.7±1.9)				11,2,7	6
2334394+193304		12.769	0.695	1.128	M8	11.2±0.2	21.0±1.8					6

Table 5—Continued

2MASSI Designation	Other Names	J	$J - H$	$J - K_S$	Spectral Type	M_J	d (pc)	$M_{J\pi}$	d_π (pc)	π Ref.	Mult. Ref.	Discovery Ref.
2336439+215338	LP 402- 58	12.761	0.660	1.053	M7	10.7±0.3	25.4±3.0					3
2347367+270206		13.186	0.734	1.191	M9	11.5±0.1	22.1±1.5					6
2349489+122438*	LP 523- 55	12.615	0.663	1.053	M8	11.2±0.2	19.6±1.6					3
2358290+270205	LP 348- 11	12.759	0.681	1.060	M6	10.1±0.4	33.8±5.8					3

Note. — M_J and distances listed are estimates obtained using our calibration while $M_{J\pi}$ and d_π are based on trigonometric parallax data. Objects included in our preliminary luminosity function ($ST \geq M7$ and $d \leq 20$ pc) are marked with *; binaries are marked with **. For multiple systems, the M_J and distance estimate (marked with parentheses) are based on the system's combined spectral type and photometry. A revised M_J and distance estimate correcting for multiplicity is given in a footnote for systems with a formal estimate of $d < 20$ pc. For multiple systems with a trigonometric parallax measurement, the combined photometry is listed in $M_{J\pi}$ and the component photometry is given in a footnote.

^aPhotometry listed is from Dahn et al. (2002) as the 2MASS photometry is contaminated by a meteor.

^b $M_J = 13.85$ for both components.

^cST change from M8 in (Gizis et al. 2000).

^dST change from M6 in PMSU.

^e $M_J = 11.80$ and 12.40 (Close, Siegler, Freed, & Biller 2003).

^f $M_J = 11.16$ and 12.01 (Close, Siegler, Freed, & Biller 2003). $d = 21.8$ pc.

^g $M_J = 11.09$ and 14.92 (Freed, Close, & Siegler 2003).

^h $M_J = 13.55$ and 13.95 (Bouy et al. 2003). $d = 21.8$ pc.

ⁱ $M_J = 11.32$ and 12.10 (Close, Siegler, Freed, & Biller 2003). $d = 24.9$ pc.

^jBased on photometric distance found by Tinney, Mould, & Reid (1993).

^k $M_J = 14.45$ and 15.10 (Bouy et al. 2003). $d = 25.4$ pc.

^lST change from M5.5 in PMSU.

^m $M_J = 11.16$ and 11.33 (Close, Siegler, Freed, & Biller 2003). $d = 24.4$ pc.

References. — Trigonometric Parallax (1) Tinney, Reid, Gizis, & Mould (1995); (2) van Altena, Lee J.T., & Hoffleit (1995) (3) Tinney (1996); (4) Dahn et al. (2002); (5) Gliese & Jahreiß (1991); (6) Monet et al. (1992); (7) ESA (1997)

References. — Multiplicity (1) Koerner, Kirkpatrick, McElwain, & Bonaventura (1999); (2) Close, Siegler, Freed, & Biller (2003); (3) Reid, Gizis, Kirkpatrick, & Koerner (2001); (4) Gizis et al. (2000); (5) Kirkpatrick et al. (2001); (6) Bouy et al. (2003); (7) Gizis et al. (2003); (8) Freed, Close, & Siegler (2003); (9) Martín, Brandner, & Basri (1999) (10) Close et al. (2002a); (11) Close et al. (2002b)

References. — Discovery (1) Irwin, McMahon, & Reid (1991); (2) McCarthy, Bertiau, & Treanor (1964); (3) NLTT; (4) Kirkpatrick et al. (2000); (5) Reid & Gilmore (1981); (6) Gizis et al. (2000); (7) Kirkpatrick et al. (1994); (8) Kirkpatrick et al. (1999); (9) Delfosse et al. (1997); (10) Martín et al. (1999); (11) Tinney, Mould, & Reid (1993); (12) Ruiz, Leggett, & Allard (1997)

Table 6. Late-type Dwarfs with $J < 9$

α (2000)	δ	V	J	$J - H$	$J - K_S$	ID
15 55 31.8	35 12 02	13.7	8.999	0.709	1.013	G 180-11, M4.5
16 58 25.3	13 58 10	13.1	8.859	0.575	1.122	G 139-3, M4
20 13 55.9	-00 52 53	10.5	8.206	0.764	1.101	BD-01 3925D
22 38 33.3	-15 18 02	12.2	6.577	0.612	1.010	EZ Aqr, Gl 866, M5.5

Table 7. M7–L8 Dwarfs Discovered Within 20 pc

2MASSI Designation	Other Names	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Spectral Type	M_J	d (pc)	Other Ref.
0019262+461407		12.609	0.676	1.135	2000 Oct 02	KP 2.1m	M8	11.2±0.2	19.5±1.6	
0019457+521317		12.820	0.748	1.204	2000 Oct 02	KP 2.1m	M9	11.5±0.1	18.7±1.2	
0109511–034326	LP 647- 13	11.695	0.774	1.277	2000 Sep 29	KP 2.1m	M9	11.5±0.1	11.1±0.7	1
0141032+180450		13.822	0.772	1.315	...	...	L4.5 ^a	13.3±0.5	12.6±2.7	2
0144353–071614		14.187	1.183	1.904	2002 Jan 24	CT 4m	L5	13.6±0.2	13.4±1.5	3,4
0148386–302439		12.282	0.641	1.038	2002 Jan 28	CT 1.5m	M7.5	11.0±0.2	18.4±1.8	
0213288+444445		13.512	0.740	1.269	2001 Jul 23	KP 4m	L1.5	12.1±0.2	18.7±1.4	
0251148–035245		13.082	0.821	1.429	2002 Jan 23	CT 4m	L3	12.7±0.2	12.1±1.1	2
0331302–304238	LP 888- 18	11.371	0.672	1.095	2000 Sep 30	KP 2.1m	M7.5 ^b	11.0±0.2	12.1±1.2	1
0417374–080000		12.166	0.654	1.112	2002 Jan 27	CT 1.5m	M7.5	11.0±0.2	17.4±1.7	
0423485–041403	SDSS J0423–0414	14.452	1.010	1.516	2002 Jan 25	CT 4m	L7.5 ^c	14.6±0.3	9.2±1.4	5,6,7,4
0429184–312356		10.887	0.680	1.086	2002 Jan 27	CT 1.5m	M7.5	(11.0±0.2) ^d	(9.7±0.9) ^d	
0435161–160657	LP 775- 31	10.396	0.616	1.060	2000 Sep 30	KP 2.1m	M7 ^e	10.7±0.3	8.6±1.0	1,8
0439010–235308		14.413	1.045	1.606	2002 Jan 26	CT 4m	L6.5	14.2±0.2	10.8±1.1	
0440232–053008	LP 655- 48	10.681	0.696	1.124	2000 Sep 29	KP 2.1m	M7	10.7±0.3	9.8±1.1	1,8,9
0443376+000205	SDSS 0443+0002	12.517	0.713	1.350	2002 Jan 28	CT 1.5m	M9.5	11.6±0.1	15.3±1.0	7,2
0445538–304820		13.409	0.835	1.425	2002 Jan 25	CT 4m	L2	12.3±0.2	16.6±1.3	
0517376–334902		11.995	0.672	1.176	2002 Jan 25	CT 4m	M8	11.2±0.2	14.7±1.2	9
0523382–140302		13.117	0.896	1.486	2002 Jan 24	CT 4m	L2.5	12.5±0.2	13.4±1.1	2
0652307+471034		13.545	1.175	1.858	2002 Jan 22	KP 4m	L4.5 ^f	13.3±0.2	11.1±1.2	
0752239+161215	LP 423- 31	10.831	0.639	1.012	2002 Jan 23	KP 4m	M7 ^g	10.7±0.3	10.5±1.2	
0835425–081923		13.149	1.195	1.993	2002 Jan 24	CT 4m	L5	13.6±0.2	8.3±0.9	
0847287–153237		13.519	0.892	1.465	2002 Jan 23	CT 4m	L2	12.3±0.2	17.5±1.4	
0859254–194926		15.505	1.067	1.778	2002 Jan 24	CT 4m	L6? ^h	14.0±0.9	19.8±8.3	
0908380+503208		14.564	1.098	1.646	2002 Jan 22	KP 4m	L5	13.6±0.2	15.9±1.8	
1006319–165326	LP 789- 23	12.041	0.620	1.041	2002 Jan 26	CT 4m	M7.5	11.0±0.2	16.4±1.6	9
1010148–040649		15.503	1.108	1.908	2002 Jan 23	CT 4m	L6	14.0±0.2	19.8±2.2	
1045240–014957	SDSS J1045–0149	13.129	0.759	1.319	2002 Jan 31	CT 1.5m	L1	12.0±0.1	16.8±1.1	7,10
1104012+195921		14.462	0.984	1.486	2002 Jan 22	KP 4m	L4	13.1±0.2	18.8±2.0	
1124048+380805		12.710	0.682	1.138	2002 Jan 25	KP 4m	M8.5	11.3±0.2	19.0±1.4	
1213033–043243		14.672	0.995	1.669	2002 Jan 24	CT 4m	L5	13.6±0.2	16.7±1.8	
1309218–233035	CE 303	11.769	0.682	1.103	2002 Jan 25	CT 4m	M8	11.2±0.2	13.3±1.1	11,10
1326298–003831	SDSS 1326–0038	16.110	1.066	1.879	...	...	L8? ⁱ	14.8±0.2	18.6±2.2	12,6
1332244–044112		12.342	0.591	1.046	2002 Jan 27	CT 1.5m	M7.5	11.0±0.2	18.9±1.8	
1356414+434258	LP 220- 13	11.704	0.673	1.070	2002 Jan 24	KP 4m	M7	10.7±0.3	15.6±1.8	
1411213–211950		12.442	0.619	1.122	2002 Jan 23	CT 4m	M9	11.5±0.1	15.7±1.0	

Table 7—Continued

2MASS Designation	Other Names	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Spectral Type	M_J	d (pc)	Other Ref.
1438082+640836		12.923	0.895	1.350	2001 Jul 21	KP 4m	M9.5	11.6±0.1	18.4±1.1	
1507277−200043		11.718	0.668	1.064	2002 Jan 27	CT 1.5m	M7.5	11.0±0.2	14.2±1.4	
1515009+484739		14.060	0.991	1.495	...	...	L6.5 ^j	14.2±0.4	9.2±1.8	2
1521010+505323		11.997	0.655	1.075	2001 Jul 20	KP 4m	M7.5	11.0±0.2	16.1±1.6	
1721039+334415		13.584	0.662	1.110	2001 Jul 19	KP 4m	L3	12.7±0.2	15.2±1.4	
1807159+501531		12.963	0.814	1.356	2001 Jul 15	KP 4m	L1.5	12.1±0.2	14.6±1.1	2
1835379+325954		10.273	0.691	1.119	2001 Jul 22	KP 4m	M8.5	11.3±0.2	6.2±0.4 ^k	13,14
2037071−113756		12.284	0.665	1.024	2001 Jul 14	KP 4m	M8	11.2±0.2	16.8±1.4	
2057540−025230		13.123	0.850	1.375	2001 Jul 15	KP 4m	L1.5 ^l	12.1±0.2	15.7±1.1	
2104149−103736		13.846	0.887	1.491	2001 Jul 15	KP 4m	L3	12.7±0.2	17.2±1.6	
2237325+392239	G 216-7B	13.346	0.664	1.192	...	...	M9.5	11.6±0.1	22.3±1.4 ^m	15

Note. — M_J and distances listed are estimates obtained using our calibration. Data for multiple systems are based on the system's combined spectral type and photometry and are marked with parentheses.

^aST from Wilson (2002)

^bST change from M6 in Paper III

^cST T0 in NIR Geballe et al. (2002)

^dBinary with $M_J = 10.96$ and 12.06 (L. Close, in preparation). $d = 11.1$ pc.

^eST change from M6 in Paper III

^fHas Li I absorption. See § 5.2

^gHas strong H α emission. See § 5.3

^hLow galactic latitude and is either a late-L dwarf or a reddened background star.

ⁱST from Fan et al. (2000)

^jST from Wilson (2002)

^k $d_\pi = 5.67 \pm 0.02$ (Paper IV)

^lHas Li I absorption and H α emission. See § 5.2

^m $d_\pi = 18.9 \pm 0.7$ (ESA 1997)

References. — (1) Paper III; (2) Wilson (2002); (3) Liebert et al. (2003); (4) Kendall et al. (2003) (5) Schneider et al. (2002); (6) Geballe et al. (2002); (7) Hawley et al. (2002); (8) McCaughrean, Scholz, & Lodieu (2002); (9) Phan-Bao et al. (2003); (10) Gizis (2002); (11) Ruiz, Wischnjewsky, Rojo, & Gonzalez (2001); (12) Fan et al. (2000); (13) Paper IV; (14) Lépine, Rich, & Shara (2003); (15) Kirkpatrick et al. (2001)

Table 8. M7–L8 Dwarfs Discovered Outside 20 pc

2MASS Designation	Other Names	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Spectral Type	M_J	d (pc)	Other Ref.
0021508+422050		14.283	0.774	1.310	2002 Jan 25	KP 4m	M9?	11.5±0.3	36.6±4.7	
0022137+120305		13.469	0.661	1.023	2001 Jul 21	KP 4m	M7.5	11.0±0.2	31.7±3.1	
0025576+391136		13.565	0.654	1.084	2002 Jan 25	KP 4m	M8	11.2±0.2	30.3±2.5	
0036448−203131		14.383	0.710	1.309	2002 Jan 25	CT 4m	M9?	11.5±0.3	38.3±4.9	
0049267−063546		13.347	0.779	1.246	2002 Jan 26	CT 4m	M8.5	11.3±0.2	25.4±1.8	
0055294+511536		13.465	0.738	1.167	2001 Jul 16	KP 4m	M7.5	11.0±0.2	31.6±3.1	
0107042+243527		13.507	0.770	1.288	2001 Jul 14	KP 4m	M9	11.5±0.1	25.6±1.7	1
0117474−340325		15.184	0.970	1.700	2002 Jan 25	CT 4m	L2?	12.3±0.3	37.6±5.9	
0120491−074103		12.976	0.684	1.122	2002 Jan 28	CT 1.5m	M8?	11.2±0.4	23.1±3.8	2
0124598+284758		13.354	0.738	1.117	2001 Jul 21	KP 4m	M8	11.2±0.2	27.5±2.3	
0141148−241731		13.437	0.673	1.135	2002 Jan 31	CT 1.5m	M7.5	11.0±0.2	31.2±2.0	
0211508+472830		12.841	0.632	1.017	2001 Jul 14	KP 4m	M7	10.7±0.3	26.4±3.1	
0214002+424336		14.239	0.780	1.277	2002 Jan 25	KP 4m	M9?	11.5±0.3	35.9±4.6	
0215159+434732	[LE 36] 2	13.301	0.744	1.111	2000 Oct 02	KP 2.1m	M7	10.7±0.3	32.6±3.8	3
0218291−313322		14.715	0.906	1.539	2002 Jan 24	CT 4m	L3	12.7±0.2	25.6±2.4	
0218578−061749	LP 649- 93	12.920	0.734	1.060	2000 Oct 01	KP 2.1m	M8 ^a	11.2±0.2	22.5±1.9	4,2
0219280−193841		14.092	0.793	1.262	2002 Jan 25	CT 4m	L1	12.0±0.1	26.2±1.8	5
0228110+253738		13.851	0.871	1.400	2002 Jan 23	KP 4m	L0?	11.7±0.3	26.6±3.2	1
0228330+181109		13.226	0.654	1.023	2002 Jan 22	KP 4m	M7	10.7±0.3	31.5±3.7	
0239424−173547		14.310	0.789	1.305	2002 Jan 26	CT 4m	L0	11.7±0.1	32.8±2.1	
0241536−124106		15.662	1.054	1.759	2002 Jan 26	CT 4m	L2?	12.3±0.3	46.8±7.4	
0257545+411132		13.964	0.754	1.237	2002 Jan 25	KP 4m	M9?	11.5±0.3	31.6±4.1	
0314401−045031		12.656	0.653	1.035	2002 Jan 23	CT 4m	M7.5	11.0±0.2	21.8±2.1	
0316451−284852		14.586	0.837	1.478	2002 Jan 24	CT 4m	L0?	11.7±0.3	37.3±4.5	
0320283−044635		13.249	0.708	1.137	2002 Jan 31	CT 1.5m	M8?	11.2±0.4	26.2±4.3	1
0326422−210205		16.111	1.337	2.226	2002 Jan 24	CT 4m	L5?	13.6±0.9	32.5±14.2	
0355047−103241		13.085	0.629	1.142	2002 Jan 25	KP 4m	M8.5	11.3±0.2	22.5±1.6	
0408290−145033		14.211	0.887	1.431	2002 Jan 24	CT 4m	L2	12.3±0.2	24.0±1.9	1
0423532−000658		13.664	0.681	1.166	2002 Jan 26	CT 4m	M8.5	11.3±0.2	29.4±2.1	
0428509−225322		13.579	0.882	1.459	...	...	L0.5 ^b	11.9±0.1	22.0±1.4	6
0430515−084900		12.954	0.689	1.174	2002 Jan 27	CT 1.5m	M8	11.2±0.2	22.9±1.9	
0436276+115124		13.788	0.699	1.203	2002 Jan 26	CT 4m	M9?	11.5±0.3	29.1±3.8	
0436278−411446		13.105	0.689	1.043	...	...	M7.5 ^c	11.0 ^c	26.6 ^c	2
0445111−060252		13.306	0.729	1.068	2002 Jan 30	CT 1.5m	M7	10.7±0.3	32.7±3.8	
0445323−364225		13.286	0.593	1.071	2002 Jan 30	CT 1.5m	M9?	11.5±0.3	23.1±3.0	
0451009−340214		13.564	0.724	1.270	2002 Jan 26	CT 4m	L0.5	11.9±0.1	21.9±1.4	
0453264−175154		15.153	1.107	1.686	2002 Jan 23	CT 4m	L3?	12.7±0.4	31.3±5.7	
0455326−270149		14.434	0.803	1.343	2002 Jan 26	CT 4m	M9?	11.5±0.3	39.2±5.1	
0508494−164716		13.713	0.749	1.213	2002 Jan 25	CT 4m	M8	11.2±0.2	32.5±2.7	
0512063−294954		15.517	1.387	2.231	2002 Jan 23	CT 4m	L4?	13.1±0.9	30.5±12.5	
0528443−325222		13.736	0.725	1.109	2002 Jan 26	CT 4m	M8.5	11.3±0.2	30.4±2.2	
0600337−331426		13.203	0.746	1.203	2002 Jan 25	CT 4m	M7.5	11.0±0.2	28.0±2.7	
0608023−294459		13.839	0.684	1.155	2002 Jan 26	CT 4m	M8.5	11.3±0.2	31.9±2.3	
0614528+453655		13.003	0.755	1.234	2002 Jan 22	KP 4m	M9	11.5±0.1	20.3±1.3	
0644143−284141		13.829	0.732	1.136	2002 Jan 25	CT 4m	M8	11.2±0.2	24.3±2.8	
0657254−401913		12.747	0.593	1.070	2002 Jan 27	CT 1.5m	M7.5	11.0±0.2	22.7±2.2	
0657557+402942		13.285	0.613	1.020	2002 Jan 23	KP 4m	M8	11.2±0.2	26.7±2.2	
0703269+463216		13.494	0.732	1.152	2002 Jan 23	KP 4m	M8	11.2±0.2	29.4±2.4	
0704493+505155		13.689	0.665	1.123	2002 Jan 23	KP 4m	M7.5	11.0±0.2	35.1±3.4	
0706285+385824		12.858	0.650	1.035	2002 Jan 25	KP 4m	M7	10.7±0.3	26.6±3.1	

Table 8—Continued

2MASS Designation	Other Names	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Spectral Type	M_J	d (pc)	Other Ref.
0710490+280909		13.398	0.662	1.040	2002 Jan 24	KP 4m	M7?	10.7 ± 0.5	34.1 ± 7.9	
0721462+193744		12.958	0.595	1.020	2002 Jan 23	KP 4m	M7.5	11.0 ± 0.2	25.1 ± 2.4	
0730489+155312		14.036	0.702	1.234	2002 Jan 22	KP 4m	M9	11.5 ± 0.1	32.7 ± 2.2	
0736017+204048		13.647	0.676	1.068	2002 Jan 25	KP 4m	M7	10.7 ± 0.3	38.3 ± 4.5	
0740557+411409		13.521	0.688	1.090	2002 Jan 23	KP 4m	M8	11.2 ± 0.2	29.7 ± 2.5	
0754054+160317		13.679	0.577	1.075	2002 Jan 23	KP 4m	M8	11.2 ± 0.2	32.0 ± 2.7	
0819460+165853		13.798	0.696	1.169	2002 Jan 23	KP 4m	M9	11.5 ± 0.1	29.3 ± 1.9	
0827202+450204		13.370	0.623	1.053	2002 Jan 24	KP 4m	M8	11.2 ± 0.2	27.7 ± 2.3	
0829324-023854		13.921	0.703	1.144	2002 Jan 25	CT 4m	M8	11.2 ± 0.2	35.7 ± 3.0	
0829490-001224		13.524	0.702	1.067	2002 Jan 31	CT 1.5m	M7?	10.7 ± 0.5	36.1 ± 8.4	
0839160+125354		13.705	0.620	1.166	2002 Jan 25	KP 4m	M9	11.5 ± 0.1	28.1 ± 1.9	
0850017-192418		12.816	0.682	1.189	2002 Jan 23	CT 4m	M8	11.2 ± 0.2	21.5 ± 1.8	
0856479+223518		15.647	1.068	1.723	2002 Jan 22	KP 4m	L3?	(12.7 ± 0.4)	(39.3 ± 7.2)	7
0857278-033239		14.378	0.804	1.349	2002 Jan 23	CT 4m	M9.5	11.6 ± 0.1	35.9 ± 2.3	
0902146-064209		13.763	0.694	1.092	2002 Jan 26	CT 4m	M7	10.7 ± 0.3	40.3 ± 4.7	
0903351-063733		13.702	0.668	1.071	2002 Jan 26	CT 4m	M7	10.7 ± 0.3	39.2 ± 4.6	
0912452+242549		13.520	0.654	1.062	2002 Jan 25	KP 4m	M7	10.7 ± 0.3	36.1 ± 4.2	
0913044-073304		13.403	0.790	1.272	2002 Jan 23	CT 4m	M9	11.5 ± 0.1	24.4 ± 1.6	
0916150+213951		13.218	0.718	1.147	2002 Jan 24	KP 4m	M8?	11.2 ± 0.7	25.9 ± 8.4	
0928256+423054		13.090	0.708	1.119	2002 Jan 24	KP 4m	M8.5	11.3 ± 0.2	22.6 ± 1.6	1
0934292-135243		13.053	0.614	1.017	2002 Jan 30	CT 1.5m	M7?	10.7 ± 0.5	29.1 ± 6.8	
0939145+395021		13.724	0.689	1.194	2002 Jan 24	KP 4m	M8?	11.2 ± 0.4	32.6 ± 5.3	
0940161+401736		13.583	0.685	1.146	2002 Jan 23	KP 4m	M7.5	11.0 ± 0.2	33.4 ± 3.3	
1003191-010507	LHS 5165/DENIS-P J1003-0105	12.352	0.667	1.085	...	...	M7 ^d	10.7 ± 0.3	21.1 ± 2.5	8,9
1011002+424503	TVLM 263-71765	13.357	0.657	1.056	...	...	M7.5 ^e	11.0^e	31.3 ± 2.8^e	10
1013279+352053		14.271	0.723	1.261	2002 Jan 23	KP 4m	M9?	11.5 ± 0.3	36.4 ± 4.7	
1017075+130839		14.111	0.879	1.435	2002 Jan 23	KP 4m	L2?	(12.3 ± 0.3)	(22.9 ± 3.6)	1,11
1018431-162427		13.771	0.643	1.117	2002 Jan 24	CT 4m	M7.5	11.0 ± 0.2	36.4 ± 3.6	
1019568+732408		12.884	0.637	1.100	2002 Jan 22	KP 4m	M8.5	11.3 ± 0.2	20.6 ± 1.5	
1028307+740841		12.911	0.644	1.009	2002 Jan 25	KP 4m	M7	10.7 ± 0.3	27.3 ± 3.2	
1054416+121408		12.483	0.670	1.020	2002 Jan 25	KP 4m	M7.5	11.0 ± 0.2	20.1 ± 2.0	
1059513-211308		14.536	0.784	1.366	2002 Jan 23	CT 4m	L1	12.0 ± 0.1	32.1 ± 2.2	
1117369+360936		14.220	0.734	1.240	2002 Jan 24	KP 4m	L0?	11.7 ± 0.3	31.5 ± 3.8	
1118387+233948		14.045	0.751	1.246	2002 Jan 24	KP 4m	M9	11.5 ± 0.1	32.8 ± 2.2	
1123360+124122		14.030	0.778	1.243	2002 Jan 25	KP 4m	M7	10.7 ± 0.3	45.6 ± 5.4	
1124552+231522		13.715	0.670	1.123	2002 Jan 25	KP 4m	M8	11.2 ± 0.2	32.5 ± 2.7	
1130476-221033		13.836	0.726	1.115	2002 Jan 24	CT 4m	M8	11.2 ± 0.2	34.4 ± 2.9	
1141440-223215		12.651	0.657	1.090	2002 Jan 23	CT 4m	M7.5	11.0 ± 0.2	21.8 ± 2.1	2
1152426+243807		13.033	0.766	1.286	2002 Jan 24	KP 4m	M9	11.5 ± 0.1	20.6 ± 1.4	
1158027-254536		13.528	0.678	1.052	2002 Jan 26	CT 4m	M8	11.2 ± 0.2	29.8 ± 2.5	
1158248+135445		13.944	0.747	1.208	2002 Jan 25	KP 4m	M9	11.5 ± 0.1	31.3 ± 2.1	
1202256-062902		13.695	0.691	1.109	2002 Jan 26	CT 4m	M9	11.5 ± 0.1	27.9 ± 1.9	
1202366-060405		13.963	0.690	1.155	2002 Jan 23	CT 4m	M8	11.2 ± 0.2	36.4 ± 3.0	
1204303+321259		13.883	0.805	1.342	2002 Jan 23	KP 4m	L0	11.7 ± 0.1	26.9 ± 1.7	1
1217293+003532		13.121	0.597	1.004	2002 Jan 24	KP 4m	M7.5	11.0 ± 0.2	27.0 ± 2.6	
1218595-055028		14.059	0.700	1.322	2002 Jan 23	CT 4m	M8.5	11.3 ± 0.2	35.3 ± 2.6	
1220116+331538		13.423	0.649	1.054	2001 Jul 20	KP 4m	M7	10.7 ± 0.3	34.5 ± 4.0	
1227154-063645		14.142	0.746	1.235	2002 Jan 24	CT 4m	M9	11.5 ± 0.1	34.3 ± 2.3	
1247357-121951		13.886	0.689	1.166	2002 Jan 24	CT 4m	M8.5	11.3 ± 0.2	32.6 ± 2.4	
1304075+403615		13.205	0.672	1.057	2001 Jul 16	KP 4m	M7	10.7 ± 0.3	31.2 ± 3.6	

Table 8—Continued

2MASS I Designation	Other Names	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Spectral Type	M_J	d (pc)	Other Ref.
1305213+224502		13.983	0.748	1.250	2002 Jan 25	KP 4m	M8	11.2±0.2	36.8±3.0	
1305410+204639		15.232	1.149	1.760	2002 Jan 24	KP 4m	L4?	13.1±0.4	26.8±5.5	
1330023−045320		13.315	0.616	1.099	2002 Jan 25	CT 4m	M8	11.2±0.2	27.0±2.2	
1339265−175505		13.424	0.706	1.134	2002 Jan 25	CT 4m	M7.5	11.0±0.2	31.1±3.0	
1413598−045748		13.394	0.695	1.161	2002 Jan 25	CT 4m	M8	11.2±0.2	28.0±2.3	
1421187−161820		12.769	0.660	1.094	2002 Jan 31	CT 1.5m	M7.5	11.0±0.2	23.0±2.2	9
1430435+291540		14.279	0.859	1.533	2002 Jan 25	KP 4m	L2	(12.3±0.2)	(24.8±2.0)	11
1436097+290035		13.312	0.665	1.142	2002 Jan 24	KP 4m	M8.5	11.3±0.2	25.0±1.8	
1438454+555913		13.095	0.684	1.058	2001 Jul 21	KP 4m	M7	10.7±0.3	29.7±3.5	
1452184+482621		13.342	0.675	1.038	2001 Jul 21	KP 4m	M7	10.7±0.3	33.2±3.9	
1456014−274735		13.269	0.611	1.066	2002 Jan 26	CT 4m	M9	11.5±0.1	22.9±1.5	
1614155+821132		13.619	0.778	1.179	2001 Jul 21	KP 4m	M8?	11.2±0.4	31.1±5.1	
1626569+395448		13.275	0.637	1.016	2001 Jul 21	KP 4m	M7.5	11.0±0.2	29.0±2.8	
1646115+501945		13.620	0.625	1.042	2001 Jul 19	KP 4m	M7	10.7±0.3	37.8±4.4	
1707333+430130		13.962	0.757	1.305	2001 Jul 19	KP 4m	L0.5 ^f	11.9±0.1	26.3±1.7	
1717140+652622	SDSS J1717+6526	14.940	1.089	1.745	...	...	L4 ^g	13.1±0.2	23.4±2.4	12
1743348+584411		14.016	0.863	1.347	2001 Jul 19	KP 4m	M9.5	11.6±0.1	30.4±1.9	
1744571+374710		13.820	0.657	1.107	2001 Jul 19	KP 4m	M7	10.7±0.3	41.4±4.8	
1801455+744229		13.135	0.666	1.098	2001 Jul 16	KP 4m	M7	10.7±0.3	30.2±3.5	
1822471+392150		13.430	0.653	1.015	2001 Jul 16	KP 4m	M7.5	11.0±0.2	31.1±3.0	
1852168+525719		13.480	0.646	1.146	2001 Jul 15	KP 4m	M8	11.2±0.2	29.2±2.4	
2004536−141622		13.169	0.660	1.117	2001 Jul 16	KP 4m	M7.5	11.0±0.2	27.6±2.7	
2014035−201621		12.527	0.665	1.092	2001 Jul 15	KP 4m	M7.5	11.0±0.2	20.5±2.0	
2019269−250244		13.698	0.714	1.249	2001 Jul 18	KP 4m	M8	11.2±0.2	32.3±2.7	
2039131−112653		13.808	0.656	1.150	2001 Jul 18	KP 4m	M8	11.2±0.2	33.9±2.8	
2047317−080820		13.670	0.704	1.078	2001 Jul 18	KP 4m	M7	10.7±0.3	38.7±4.5	
2107316−030733		14.222	0.772	1.306	2001 Jul 18	KP 4m	L0	11.7±0.1	31.5±2.0	
2214506−131959		13.458	0.744	1.133	2001 Jul 15	KP 4m	M7.5	11.0±0.2	31.5±3.1	
2228304+344034		13.538	0.633	1.025	2001 Jul 18	KP 4m	M7	10.7±0.3	36.4±4.2	
2238074+435317		13.839	0.809	1.320	2001 Jul 18	KP 4m	L1.5	12.1±0.2	21.8±1.6	
2247017+195528		13.829	0.738	1.224	2001 Jul 19	KP 4m	M9	11.5±0.1	29.7±2.0	
2252014−181558		13.570	0.730	1.158	2001 Jul 19	KP 4m	M8.5	11.3±0.2	28.2±2.0	
2254519−284025		14.161	0.717	1.264	2001 Jul 18	KP 4m	L0.5	11.9±0.1	28.8±1.9	
2259324+445028		13.340	0.598	1.006	2001 Jul 15	KP 4m	M7.5	11.0±0.2	29.9±2.9	
2326266+134552		13.630	0.621	1.044	2001 Jul 19	KP 4m	M7.5	11.0±0.2	34.1±3.4	
2329129+270415		13.593	0.708	1.065	2001 Jul 21	KP 4m	M7	10.7±0.3	37.3±4.4	
2336164+183500		13.650	0.712	1.196	2001 Jul 21	KP 4m	M8.5	11.3±0.2	29.2±2.1	
2337149−083808	LP 763- 38	12.246	0.643	1.040	2001 Jul 21	KP 4m	M7	10.7±0.3	20.1±2.4	4
2351296+451926		13.492	0.651	1.025	2001 Jul 15	KP 4m	M7 ^h	10.7±0.3	35.6±4.2	

Note. — M_J and distances listed are estimates obtained using our calibration. Data for multiple systems are based on the system's combined spectral type and photometry and are marked with parentheses.

^aST change from M6 in Paper III

^bST from Kendall et al. (2003)

^cData listed are based on photometric distance from Phan-Bao et al. (2003).

^dST from Gizis (2002)

^eData listed are based on photometric distance from Tinney, Reid, Gizis, & Mould (1995).

^fDisplays strong H α emission. See § 5.3

^gST from Hawley et al. (2002)

^hDisplays strong H α emission. See § 5.3

References. — (1) Wilson (2002); (2) Phan-Bao et al. (2003); (3) Luyten & Ebbighausen (1936); (4) Paper III; (5) Lodieu, Scholz, & McCaughrean (2002); (6) Kendall et al. (2003); (7) Bouy et al. (2003); (8) Phan-Bao et al. (2001); (9) Gizis (2002); (10) Tinney, Reid, Gizis, & Mould (1995); (11) Gizis et al. (2003); (12) Hawley et al. (2002)

Table 9. Early-type M Dwarfs Discovered Within 20 pc

2MASSJ Designation	Other Names	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Spectral Type	M_J	d (pc)	Other Ref.
0035441–054110	LP 645- 53	10.717	0.633	1.001	2000 Sep 30	KP 2.1m	M5	9.4±0.3	18.7±2.3	1
0214125–035743	LHS 1363/LP 649- 72	10.472	0.633	1.006	2000 Sep 30	KP 2.1m	M6.5 ^a	10.5±0.3	10.1±1.4	1
0334106–213034		11.957	0.706	1.121	2002 Jan 27	CT 1.5m	M6	10.1±0.4	23.3±4.0	
0354200–143738		11.358	0.602	1.033	2002 Jan 28	CT 1.5m	M6.5	10.5±0.3	15.2±2.1	
0410480–125142	LP 714- 37	11.060	0.591	1.045	2000 Sep 30	KP 2.1m	M6 ^b	10.1±0.4	15.4±2.7	1,2,3
1124532+132253		11.042	0.643	1.014	2002 Jan 24	KP 4m	M5.5	9.6±0.2	19.8±1.9	
1424187–351432		11.809	0.625	1.019	2002 Jan 27	CT 1.5m	M6.5	10.5±0.3	18.6±2.6	
1431304+171758		12.217	0.649	1.054	2002 Jan 24	KP 4m	M6.5	10.5±0.3	22.5±3.2	
1847034+552243		11.919	0.641	1.021	2001 Jul 16	KP 4m	M6.5	10.5±0.3	19.6±2.8	
2132297–051158	LP 698- 2	11.439	0.724	1.054	2001 Jul 15	KP 4m	M6 ^c	10.1±0.4	18.4±3.2	1,3

^aST change from M5.5 in Paper III^bST change from M5.5 in Paper III^cST change from M5.5 in Paper III

References. — (1) Paper III; (2) Wilson (2002); (3) Phan-Bao et al. (2003)

Table 10. Early-type M dwarfs Discovered Outside 20 pc

2MASS Designation	Other Names	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Spectral Type	M_J	d (pc)	Other Ref.
0014083+493433		13.276	0.621	1.022	2001 Jul 15	KP 4m	M6	10.1±0.4	42.8±7.4	
0017533+145324		13.204	0.690	1.056	2001 Jul 16	KP 4m	M6.5	10.5±0.3	35.5±5.0	1
0030198+314125		13.383	0.630	1.006	2001 Jul 21	KP 4m	M4	8.9±0.2	77.4±6.2	
0104081+362301		13.064	0.637	1.015	2001 Jul 23	KP 4m	M5	9.4±9.3	54.8±7.1	
0219331+141632		12.725	0.660	1.014	2001 Jul 23	KP 4m	M4	7.6±0.3	104±13	
0234295+361311		12.459	0.637	1.002	2000 Sep 30	KP 2.1m	M6 ^a	10.1±0.4	29.4±5.1	2
0250023+080841	LP 651- 17/LHS 1450	11.878	0.652	1.028	2000 Sep 30	KP 2.1m	M5.5	9.7±0.2	26.9±2.7	2
0301032+441656		12.059	0.681	1.061	2000 Oct 02	KP 2.1m	M6	10.1±0.4	24.5±4.2	
0336489+241801		9.905	0.530	1.003	2002 Jan 27	CT 1.5m	M3	7.5±0.2	30.3±2.8	
0413398+270429	LP 890- 2/DENIS-P J0413+2704	12.214	0.636	1.024	2000 Sep 30	KP 2.1m	M6	10.1±0.4	26.3±4.5	2,3
0422205+360608		12.421	0.650	1.056	2002 Jan 27	CT 1.5m	M6.5	10.5±0.3	24.7±3.5	
0439340+323551		11.601	0.560	1.006	2002 Jan 27	CT 1.5m	M5.5	9.5±0.5	26.0±5.7	
0502386+322750		12.458	0.623	1.047	2002 Jan 26	CT 4m	M6.5	10.5±0.3	25.1±3.6	
0701058+261412		13.112	0.661	1.045	2002 Jan 24	KP 4m	M6.5	10.5±0.3	34.0±4.8	
0835432+313932		13.528	0.639	1.029	2002 Jan 24	KP 4m	M6	10.1±0.4	48.1±8.3	
0904465+474634		13.573	0.660	1.028	2002 Jan 25	KP 4m	M5	9.7±0.3	60.6±9.3	
0926332+015102		12.931	0.656	1.040	2002 Jan 26	CT 4m	M6	10.1±0.4	36.5±6.3	
0951328+241737		13.309	0.557	1.056	2002 Jan 24	KP 4m	M5	9.8±0.2	49.9±4.4	
1010426+194406		13.281	0.652	1.080	2002 Jan 24	KP 4m	M6.5	10.5±0.3	36.7±5.2	
1021513+032309		12.359	0.612	1.007	2002 Jan 27	CT 1.5m	M6.5	10.5±0.3	24.0±3.4	4
1104335+051043		13.376	0.644	1.112	2002 Jan 24	CT 4m	M6	10.1±0.4	44.8±7.7	
1129338+023311		9.539	0.542	1.052	2002 Jan 27	CT 1.5m	M2	6.9±0.2	34.3±3.2	
1231546+513039		13.243	0.649	1.005	2002 Jan 24	KP 4m	M6.5	10.5±0.3	36.1±5.1	
1233231+284158		13.254	0.678	1.051	2001 Jul 20	KP 4m	M5.5	9.5±0.3	56.9±6.6	
1242283+265251		12.649	0.632	1.002	2001 Jul 16	KP 4m	M5	9.4±0.5	44.0±9.1	
1252170+335739	LP 267-299	12.246	0.645	1.007	2001 Jul 16	KP 4m	M5.5	9.4±0.2	36.4±3.7	
1340115+145159		13.530	0.678	1.041	2002 Jan 26	CT 4m	M6.5	10.5±0.3	41.2±5.8	
1344492+122848		12.064	0.599	1.001	2002 Jan 31	CT 1.5m	M3	7.3±0.2	90.1±9.2	
1431156+131824	LP 740- 25	11.136	0.640	1.015	2002 Jan 27	CT 1.5m	M5	9.2±0.3	23.9±3.0	
1445062+440939		12.439	0.633	1.018	2001 Jul 21	KP 4m	M6.5	10.5±0.3	24.9±3.5	
1511512+303306		12.956	0.688	1.048	2001 Jul 21	KP 4m	M5	9.5±0.7	49.8±16.5	
1516221+531631		13.170	0.679	1.066	2001 Jul 21	KP 4m	M6	10.1±0.4	40.8±7.0	
1527192+413044		12.489	0.660	1.066	2001 Jul 14	KP 4m	M6.5	10.5±0.3	25.5±3.6	
1543554+531521		13.325	0.604	1.041	2001 Jul 21	KP 4m	M5	9.1±0.2	71.6±8.0	
1623388+161554		12.758	0.663	1.020	2001 Jul 19	KP 4m	M5	9.5±0.2	44.4±4.7	
1626353+251235		12.811	0.612	1.006	2001 Jul 15	KP 4m	M6.5	10.5±0.3	29.6±4.2	

Table 10—Continued

2MASS Designation	Other Names	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Spectral Type	M_J	d (pc)	Other Ref.
1626373+604043		12.647	0.687	1.071	2001 Jul 15	KP 4m	M6.5	10.5±0.3	27.4±3.9	
1732036+155715		12.494	0.636	1.040	2001 Jul 15	KP 4m	M3	7.4±0.2	104±10	
1732281+300454		13.331	0.613	1.018	2001 Jul 16	KP 4m	M6.5	10.5±0.3	37.6±5.3	
1809281+393608		13.204	0.629	1.024	2001 Jul 16	KP 4m	M5.5	9.5±0.2	55.1±5.6	
1810005+405600		11.275	0.632	1.004	2001 Jul 18	KP 4m	M3	7.3±0.2	63.2±6.8	
2015194−160133		12.713	0.661	1.021	2001 Jul 15	KP 4m	M5.5	9.8±0.5	38.8±8.4	
2033573−042941		13.509	0.654	1.012	2001 Jul 18	KP 4m	M6.5	10.5±0.3	40.8±5.8	
2041410−033353	LP 695-351	12.528	0.646	1.024	2001 Jul 16	KP 4m	M6	10.1±0.4	30.3±5.2	
2115408+165716		11.331	0.572	1.099	2000 Oct 01	KP 2.1m	M4	8.9±0.2	30.8±2.3	
2125458−001834		13.153	0.684	1.051	2001 Jul 15	KP 4m	M6.5	10.5±0.3	34.6±4.9	
2226368−023950		12.559	0.654	1.010	2001 Jul 14	KP 4m	M6.5	10.5±0.3	26.3±3.7	
2240386−025056		13.220	0.610	1.007	2001 Jul 15	KP 4m	M6.5	10.5±0.3	35.7±5.1	
2246444+294135		13.117	0.657	1.007	2001 Jul 15	KP 4m	M5.5	9.7±0.2	48.2±5.4	
2246527+225517		12.012	0.626	1.014	2001 Jul 15	KP 4m	M5	9.2±0.3	36.3±4.6	
2252051+255118		13.099	0.654	1.008	2001 Jul 15	KP 4m	M5	9.3±0.2	57.8±6.2	
2317207−023632		12.304	0.642	1.029	2000 Sep 29	KP 2.1m	M6.5 ^b	10.5±0.3	23.4±3.3	2
2337383−125027	LP 763- 3/WT 2339	11.461	0.610	1.034	2001 Jul 21	KP 4m	M6 ^c	10.1±0.4	18.6±3.2	2,4

^aST change from M5 in Paper III^bST change from M5.5 in Paper III^cST change from M5.5 in Paper III

References. — (1) Wilson (2002); (2) Paper III; (3) Phan-Bao et al. (2001); (4) Phan-Bao et al. (2003)

Table 11. Young Objects

2MASS Designation	J	$J - H$	$J - K_S$	Obs. date	Telescope	Comments
0253597+320637	13.623	0.693	1.079	2002 Jan 25	KP 4m	
0435145-141446	11.868	1.236	1.927	2002 Jan 24	CT 4m	
0608528-275358	13.600	0.702	1.210	2002 Jan 25	CT 4m	enhanced VO absorption
0619526-290359	15.101	0.882	1.649	2002 Jan 24	CT 4m	
2234416+404138	12.569	0.735	1.138	2000 Oct 02	KP 2.1m	strong $H\alpha$ emission

Table 12. Spectroscopically Confirmed Giants

2MASS Designation	J	$J - H$	$J - K_S$	Obs. date	Telescope	Spectral Type	Comments
0012342–225516	10.393	0.779	1.125	2002 Jan 31	CT 1.5m	M1 III	
0028097+121719	10.177	0.883	1.392	2000 Oct 02	KP 2.1m	M5 III	
0036441–060858	9.998	0.785	1.139	2002 Jan 28	CT 1.5m	M1 III	
0229220–110612	13.213	0.687	1.325	2002 Jan 31	CT 1.5m	M0 III	
0252543–114220	12.505	0.813	1.171	2002 Jan 30	CT 1.5m	M1 III	
0424503–740443	12.337	0.760	1.162	2002 Jan 26	CT 4m	M8 III	
0436168–641131	12.098	0.863	1.275	2002 Jan 26	CT 4m	M5 III	
0534324–121631	10.161	0.721	1.078	2002 Jan 27	CT 1.5m	M0 III	
0613001–163648	9.144	0.758	1.091	2002 Jan 27	CT 1.5m	M1 III	
0614035–230347	9.976	0.832	1.249	2002 Jan 27	CT 1.5m	M4 III	
0636411–324041	9.082	0.737	1.052	2002 Jan 27	CT 1.5m	M0 III	
0647027–622725	13.271	0.510	1.067	2002 Jan 31	CT 1.5m	M3 III	
0827311–110002	15.834	1.210	1.930	2002 Jan 25	CT 4m	M0 III	
0846564–204807	9.835	0.858	1.313	2002 Jan 27	CT 1.5m	M6 III	
1010187–043536	9.579	0.764	1.087	2002 Jan 27	CT 1.5m	M1 III	
1035438–091625	9.795	0.697	1.107	2002 Jan 27	CT 1.5m	M3 III pec	variable H α emission
1040446–080950	9.725	0.683	1.008	2002 Jan 27	CT 1.5m	M1 III	
1307008–240531	9.514	0.858	1.498	2002 Jan 27	CT 1.5m	M8 III	
1401078–004744	11.487	0.682	1.099	2002 Jan 24	KP 4m	M6 III	
1442162–300709	9.281	0.790	1.106	2002 Jan 27	CT 1.5m	M3 III	
1447033–134423	9.161	0.800	1.116	2002 Jan 27	CT 1.5m	M1 III	
1501024–240826	9.838	0.633	1.113	2002 Jan 27	CT 1.5m	M8 III	
1502046–342517	9.551	0.684	1.016	2002 Jan 27	CT 1.5m	M0 III	
1553258+530924	9.974	0.770	1.170	2001 Jul 20	KP 4m	M8 III	
1553557–220038	10.425	0.719	1.028	2002 Jan 28	CT 1.5m	M5 III	
1604434–191014	10.389	0.734	1.047	2002 Jan 28	CT 1.5m	M5 III	
2002154–031431	9.913	0.914	1.409	2001 Jul 16	KP 4m	M4 III	
2019460–013400	9.363	0.833	1.475	2001 Jul 15	KP 4m	M6 III	

Table 12—Continued

2MASSI Designation	J	$J - H$	$J - K_S$	Obs. date	Telescope	Spectral Type	Comments
2025585–274618	11.117	1.253	2.033	2001 Jul 15	KP 4m	M6 III	
2231451–233245	11.451	0.766	1.200	2001 Jul 16	KP 4m	M3 III pec	H α in emission
2244246+133701	10.120	1.169	1.829	2000 Oct 02	KP 2.1m	M4 III	
2332026+243843	11.532	0.819	1.193	2000 Oct 02	KP 2.1m	M3 III	
2336511+482441	13.471	0.907	1.339	2001 Jul 14	KP 4m	M5 III	
2346447+160359	11.502	0.917	1.346	2000 Oct 02	KP 2.1m	M7 III	

Table 13. Spectroscopically Confirmed Carbon Stars

2MASS Designation	J	$J - H$	$J - K_S$	Obs. Date	Telescope	Visual pm ^a (epochs)	Comments
0236246–204103	13.673	0.768	1.224	2002 Jan 26	CT 4m	yes (12/79–11/98)	dC, LP 830-18 ^b
0402332–681623	13.490	1.420	2.372	2002 Jan 31	CT 1.5m	no (12/92–01/00)	
0426387+142516	13.227	1.211	2.014	2000 Oct 02	KP 2.1m	no (12/55–11/98)	
0431342–723208	13.185	1.234	1.958	2002 Jan 31	CT 1.5m	no (01/93–12/98)	
0459495–750918	12.638	1.327	2.325	2002 Jan 26	CT 4m	no (01/75–12/98)	
0614383–135137	9.829	0.789	1.127	2002 Jan 27	CT 1.5m	no (11/83–01/99)	
0635542–645226	12.826	0.945	1.395	2002 Jan 23	CT 4m	no (11/82–02/00)	
0654386–713059	13.060	1.244	2.083	2002 Jan 30	CT 1.5m	no (02/91–12/98)	
0701081–681854	12.816	1.307	2.273	2002 Jan 30	CT 1.5m	no (02/78–12/98)	
0837170–132114	9.250	1.046	1.548	2002 Jan 27	CT 1.5m	no (01/82–02/99)	
0843554–122407	9.532	0.991	1.511	2002 Jan 27	CT 1.5m	no (01/84–03/98)	
1501069–053138	13.571	1.223	2.065	2002 Jan 31	CT 1.5m	no (07/84–02/99)	
1515110–133227	12.594	1.078	1.809	2002 Jan 31	CT 1.5m	no (04/82–03/98)	Gizis (2002)
1622328+423753	13.307	0.776	1.385	2001 Jul 15	KP 4m	yes (05/53–06/98)	dC, LP 225-12 ^b
2206536–250628	10.934	1.178	2.012	2001 Jul 14	KP 4m	no (07/77–11/98)	
2217099–260703	11.056	1.233	2.174	2001 Jul 14	KP 4m	no (06/80–07/98)	

^aPossible visual proper motion determined by P.J.L. from examining DSS, XDSS and 2MASS images. Epochs listed are either DSS–2MASS or XDSS R –2MASS.

^bA discussion of these new carbon dwarfs will appear in P.J. Lowrance et al., in preparation.

Table 14. M_J /Spectral Type Calibration Data

Name	Spectral Type ^a	M_J	Ref.
Gl 551	5.5	9.68 ± 0.01	1
GJ 1286	5.5	9.90 ± 0.05	1
GJ 1002	5.5	9.98 ± 0.03	1
LHS 3339	5.5	10.05 ± 0.06	3
Gl 412B	6	10.04 ± 0.01	1
LHS 1443	6	10.09 ± 0.20	2
LHS 1516	6	10.11 ± 0.26	2
Gl 406	6	10.15 ± 0.01	1
GJ 1111	6.5	10.40 ± 0.02	1
LHS 191	6.5	10.43 ± 0.08	3
LHS 2471	6.5	10.47 ± 0.09	3
LHS 523	6.5	10.60 ± 0.11	2
LHS 292	6.5	10.62 ± 0.03	1
LHS 2930	6.5	10.77 ± 0.04	3
LHS 429	7	10.72 ± 0.03	3
LHS 3003	7	10.91 ± 0.06	3
GRH 2208-20	7.5	10.88 ± 0.07	3
TVLM 832-10443	8	10.83 ± 0.04	3
LP 412-31	8	10.98 ± 0.05	3
LHS 2397a	8	11.13 ± 0.08	3
BRI 0246-1703	8	11.42 ± 0.19	3
GL 569Ba	8.5	11.18 ± 0.08	3
CTI 012657.5+280202	8.5	11.44 ± 0.06	3
TVLM 513-46546	8.5	11.72 ± 0.06	3
BRI 1222-1222	9	11.34 ± 0.14	3
TVLM 868-110639	9	11.49 ± 0.17	3
LHS 2065	9	11.55 ± 0.04	3
LHS 2924	9	11.66 ± 0.04	3
GL 569Bb	9	11.69 ± 0.08	3
BRI 0021-0214	9.5	11.44 ± 0.11	3
2MASSW J0149090+295613	9.5	11.63 ± 0.05	3
PC 0025+0447	9.5	11.78 ± 0.25	3

Table 14—Continued

Name	Spectral Type ^a	M_J	Ref.
2MASSP J0345432+254023	10	11.77 ± 0.06	3
HD 89744B	10	11.93 ± 0.07	3
2MASSW J0746425+200032A	10.5	11.81 ± 0.05	3
2MASSW J1439284+192915	11	11.87 ± 0.04	3
2MASSW J1658038+702702	11	11.97 ± 0.04	3
GJ 1048B	11	12.03 ± 0.13	3
Kelu 1	12	12.03 ± 0.09	3
GL 618.1B	12.5	12.90 ± 0.18	3
2MASSW J1146345+223053A	13	12.58 ± 0.08	3
DENIS-P 1058.7–1548	13	12.95 ± 0.06	3
2MASSW J0036159+182110	13.5	12.71 ± 0.04	3
2MASSW J0326137+295015	13.5	12.90 ± 0.12	3
GD 165B	14	13.23 ± 0.17	3
2MASSW J1112257+354813	14.5	12.88 ± 0.06	3
2MASSW J2224438–015852	14.5	13.77 ± 0.04	3
LHS 102B	15	13.20 ± 0.24	3
2MASSW J1328550+211449	15	13.38 ± 0.27	3
2MASSW J1507476–162738	15	13.49 ± 0.04	3
DENIS-P 1228.2–1547	15	13.56 ± 0.10	3
DENIS-P 0205.4–1159	17	13.85 ± 0.08	3
2MASSI J0825196+211552	17.5	14.97 ± 0.04	3
GL 337C	18	14.14 ± 0.09	3
2MASSW J1523226+301456	18	14.79 ± 0.08	3
2MASSW J1632291+190441	18	14.89 ± 0.09	3

^aM5 = 5, L0 = 10, L8 = 18

References. — (1) 8 pc Sample; (2) PMSU; (3) Dahn et al. (2002)

Table 15. $J < 9$ Sources with IRAS Catalog Counterparts

α (2000)	δ	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
00 01 21.4	51 12 13	0.920	0.700	6.910	1.620	IRAS 23587+5055
00 20 53.4	50 09 07	0.930	0.590	5.480	1.520	IRAS 00182+4952
00 24 18.7	34 23 04	...	...	4.830	1.060	IRAS 00216+3406, M0
01 49 43.1	38 40 53	...	...	4.960	1.100	IRAS 01467+3826
02 39 05.1	31 00 26	0.850	0.450	6.070	1.300	IRAS 02361+3047
03 03 52.0	45 11 42	0.950	0.530	5.360	1.480	IRAS 03005+4459
03 06 47.1	28 20 32	6.300	...	4.890	1.410	IRAS 03037+2808
03 15 24.2	40 44 54	0.800	0.320	5.350	1.120	IRAS 03121+4033
04 02 39.4	21 03 59	0.770	0.510	5.120	1.280	IRAS 03597+2055
04 17 46.0	18 22 36	0.990	0.520	5.840	1.510	IRAS 04148+1815
04 24 01.0	-67 07 01	0.869	0.463	5.460	1.332	IRAS 04238-6713
05 37 50.0	-15 48 11	0.984	0.690	4.890	1.674	IRAS 05355-1549
05 47 58.6	-33 05 10	0.913	0.637	6.167	1.550	IRAS 05461-3306
05 48 11.9	-32 00 10	0.846	0.453	5.202	1.299	IRAS 05463-3201
05 49 06.1	-22 41 14	0.786	0.412	6.490	1.198	IRAS 05469-2242
06 09 13.1	-27 43 09	1.263	0.931	6.308	2.194	IRAS 06072-2742
06 10 52.7	-14 59 44	0.976	0.482	6.470	1.458	IRAS 06086-1459
06 24 21.2	-18 30 15	1.075	0.584	6.535	1.659	IRAS 06222-1828
06 25 52.9	-31 22 06	0.915	0.571	6.756	1.486	IRAS 06329-3120
06 32 49.5	-63 35 49	0.780	0.604	5.607	1.384	IRAS 06325-6333
06 38 51.6	-24 23 34	0.825	0.415	6.410	1.240	IRAS 06367-2420
06 40 27.8	-27 41 22	0.834	0.373	5.836	1.207	IRAS 06384-2738
06 41 54.8	51 29 15	0.893	0.521	4.949	1.414	IRAS 06379+5132
06 50 35.9	-29 59 22	0.942	0.606	6.088	1.548	IRAS 06486-2955
06 51 27.8	-36 26 32	0.776	0.620	6.007	1.396	IRAS 06497-3622
06 51 38.3	-34 55 57	0.867	0.514	5.271	1.381	IRAS 06498-3452
06 51 40.4	-59 55 18	0.848	0.437	6.264	1.285	IRAS 06509-5951
06 55 12.4	-36 07 10	0.830	0.432	6.098	1.262	IRAS 06534-3603
06 58 07.2	-38 37 53	0.854	0.558	6.307	1.412	IRAS 06564-3833
07 00 16.2	-34 51 16	0.895	0.538	5.120	1.433	IRAS 06584-3447
07 03 09.0	-36 53 19	0.963	0.615	6.009	1.578	IRAS 07013-3648
07 06 59.3	-37 55 50	0.826	0.474	6.359	1.300	IRAS 07052-3751
07 25 18.3	16 58 00	0.775	0.345	5.371	1.120	IRAS 07224+1703
07 39 07.8	44 27 43	6.18:	...	5.069	1.110	IRAS 07355+4434
08 09 19.4	39 39 07	0.773	0.350	5.237	1.123	IRAS 08059+3947
08 10 29.3	24 01 47	0.906	0.466	4.990	1.372	IRAS 08074+2410
08 15 11.1	-04 27 50	1.083	0.687	6.641	1.770	IRAS 08126-0418
08 15 52.4	-01 50 49	0.937	0.470	5.094	1.407	IRAS 08133-0141
08 26 26.8	-03 17 43	0.847	0.481	5.838	1.328	IRAS 08239-0307
09 42 10.0	-14 09 16	0.819	0.477	5.075	1.296	IRAS 09397-1355
10 36 17.3	59 56 09	6.00:	...	4.714	1.285	IRAS 10330+6011
12 14 52.7	-10 02 31	0.796	0.333	5.527	1.129	IRAS 12123-0946
13 25 31.7	-45 13 49	0.825	0.460	6.380	1.285	IRAS 13225-4458
13 29 16.5	-33 37 35	0.828	0.451	6.524	1.279	IRAS 13264-3321
13 29 38.9	-23 45 02	0.901	0.518	6.287	1.419	IRAS 13269-2329
13 48 44.6	33 43 34	0.788	0.448	5.482	1.236	IRAS 13465+3358
14 33 28.3	17 36 46	0.773	0.534	6.841	1.307	IRAS 14311+1749, CO Boo
14 35 48.9	-23 36 29	0.788	0.619	5.145	1.407	IRAS 14329-2323
14 46 44.7	-26 45 13	0.909	0.469	5.988	1.378	IRAS 14438-2632

Table 15—Continued

α (2000)	δ	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
14 56 01.6	−26 42 38	0.836	0.525	5.902	1.361	IRAS 14530-2630
15 02 37.6	−27 50 32	0.757	0.505	5.329	1.262	IRAS 14596-2738
15 04 27.5	−30 52 46	0.784	0.566	6.968	1.350	IRAS 15014-3041
15 07 28.1	−35 05 35	0.934	0.572	5.516	1.506	IRAS 15043-3454
15 15 07.5	−19 41 06	0.888	0.450	5.511	1.338	IRAS 15121-1929
15 26 20.8	−26 32 08	0.829	0.637	6.934	1.466	IRAS 15233-2621
15 45 33.6	−20 27 42	0.924	0.741	5.465	1.665	IRAS 15426-2018
15 53 58.9	−18 30 16	0.911	0.582	5.300	1.493	IRAS 15511-1821
16 03 44.2	−13 06 49	0.868	0.484	5.026	1.352	IRAS 16009-1258
16 05 27.5	−18 55 04	0.872	0.577	5.183	1.449	IRAS 16025-1846
16 19 02.4	59 50 11	5.91:	...	4.822	1.089	IRAS 16182+5957
17 05 27.6	32 05 45	0.790	0.313	5.347	1.103	IRAS 17035+3209
17 06 21.4	14 27 35	0.859	0.471	6.364	1.330	IRAS 17041+1431
17 17 00.1	60 46 43	0.749	0.300	5.412	1.049	IRAS 17163+6049
17 45 01.1	16 51 44	0.870	0.446	5.276	1.316	IRAS 17428+1652
17 57 57.9	15 00 58	0.934	0.523	5.608	1.457	IRAS 17556+1501
18 03 49.2	60 10 15	0.794	0.434	6.435	1.228	IRAS 18031+6009
18 09 30.7	31 12 30	0.945	0.480	7.145	1.425	IRAS 180976+3111
18 17 24.5	42 36 16	0.766	0.470	6.516	1.236	IRAS 18158+4235
18 19 30.4	35 00 21	0.841	0.423	5.409	1.264	IRAS 18177+3459
18 23 10.3	25 41 07	0.831	0.533	6.566	1.364	IRAS 18211+2539
18 23 18.0	24 43 16	0.812	0.598	7.160	1.410	IRAS 18212+2441
18 25 29.6	31 33 05	0.802	0.573	6.892	1.375	IRAS 18236+3131
18 32 53.8	43 01 03	0.847	0.624	6.441	1.471	IRAS 18313+4258
18 56 49.2	47 57 10	1.107	0.674	6.225	1.781	IRAS 18554+4753
19 21 16.8	−22 39 57	0.864	0.450	5.571	1.314	IRAS 19182-2245
19 22 05.0	−38 50 38	0.847	0.628	6.351	1.475	IRAS 19186-3856
19 34 33.4	−33 48 18	0.943	0.482	5.582	1.425	IRAS 19312-3354
19 36 44.5	−15 19 25	1.029	0.523	5.580	1.552	IRAS 19339-1526
19 37 37.7	−14 36 51	1.089	0.702	6.162	1.791	IRAS 19348-1443
19 39 06.4	−16 16 58	0.867	0.643	5.609	1.510	IRAS 19362-1623
19 39 27.7	−13 00 01	0.927	0.517	6.502	1.444	IRAS 19366-1306
19 42 08.9	−28 46 11	0.857	0.453	7.397	1.310	IRAS 19390-2853
19 42 29.4	−36 04 35	1.020	0.489	4.927	1.509	IRAS 19392-3611
19 42 34.7	−13 55 58	0.941	0.536	5.242	1.477	IRAS 19397-1403
19 43 04.8	−21 00 53	0.889	0.457	4.986	1.346	IRAS 19401-2108
19 44 59.5	−31 14 14	1.003	0.482	5.427	1.485	IRAS 19417-3121
19 49 23.1	−17 58 21	0.883	0.498	5.457	1.381	IRAS 19464-1805
19 58 28.5	−02 27 28	0.908	0.514	5.967	1.422	IRAS 19558-0235
19 58 43.2	−36 51 44	0.943	0.505	5.265	1.448	IRAS 19554-3659
19 59 58.0	−22 58 14	0.907	0.556	5.833	1.463	IRAS 19570-2306
20 07 02.2	−15 23 25	0.924	0.532	4.958	1.456	IRAS 20042-1532
20 09 17.8	−28 39 30	0.822	0.526	4.985	1.348	IRAS 20062-2848
20 13 15.9	−19 53 55	0.952	0.511	5.281	1.463	IRAS 20103-2003
20 18 44.6	−20 54 45	0.858	0.592	5.624	1.450	IRAS 20158-2104
20 21 09.3	−19 23 42	0.944	0.700	5.792	1.644	IRAS 20182-1933
20 23 53.1	−12 49 17	0.758	0.495	6.733	1.253	IRAS 20211-1258
20 27 18.5	−01 59 20	0.927	0.558	5.623	1.485	IRAS 20247-0209
20 29 23.8	00 49 02	1.006	0.492	5.105	1.498	IRAS 20268-0059

Table 15—Continued

α (2000)	δ	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
20 30 03.3	−28 48 49	0.942	0.498	5.691	1.440	IRAS 20270-2858
20 34 13.3	00 50 24	0.887	0.468	5.432	1.355	IRAS 20316-0100
21 10 17.1	18 34 40	5.950	...	4.530	1.420	IRAS 21079+1822
21 12 09.9	18 49 02	0.820	0.500	5.380	1.320	IRAS 21098+1836
21 12 16.7	22 17 55	0.960	0.660	7.290	1.620	IRAS 21100+2205
21 12 31.7	21 29 59	5.820	...	4.470	1.350	IRAS 21102+2117, M7
21 14 45.4	−27 58 41	0.818	0.663	6.824	1.481	IRAS 2117-2811
21 26 56.9	13 16 54	0.780	0.470	5.390	1.250	IRAS 21245+1303
21 35 26.6	17 24 06	0.790	0.580	7.260	1.370	IRAS 21330+1710
21 35 26.6	16 26 01	0.800	0.420	5.630	1.220	IRAS F21330+1612
21 59 04.0	−25 32 44	0.913	0.487	4.860	1.400	IRAS 21562-2547
22 34 23.9	44 28 07	5.750	...	...	...	IRAS 22322+4412
22 35 28.4	40 10 25	6.210	...	4.880	1.330	IRAS 22332+3954
22 56 00.7	43 23 39	0.850	0.550	5.790	1.400	IRAS 22537+4307
22 58 35.1	30 24 38	0.710	0.430	6.990	1.140	IRAS 22562+3008
22 59 56.8	35 09 48	0.830	0.460	5.980	1.290	IRAS 22575+3453
23 07 48.1	29 17 55	0.910	0.490	5.700	1.400	IRAS 23053+2901
23 14 06.3	31 04 42	0.840	0.370	5.560	1.210	IRAS 23116+3048
23 19 31.3	46 52 25	1.120	0.570	6.160	1.690	IRAS 23171+4635, C*

Table 16. $J < 9$ Sources with Stellar Counterparts

α (2000)	δ	V	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
00 20 01.3	51 53 26	8.72	5.630	...	...	...	HD 232159, K5
00 38 06.2	37 35 21	10.4	0.720	0.340	5.900	1.060	GSC 02784-01363, M
00 42 32.4	33 11 31	9.57	0.740	0.300	5.480	1.040	BD+32 116, K5
00 44 41.2	47 55 47	9.61	0.670	0.490	5.920	1.160	BD+47 182, K5
01 15 51.7	-20 34 59	10.1	0.689	0.325	6.185	1.014	BD-21 199
01 28 55.4	34 40 19	10.4	0.690	0.310	6.220	1.000	GSC 02300-01336
01 48 29.8	40 16 31	10.02	0.690	0.350	5.650	1.040	GSC 02819-01779
02 04 31.5	15 37 48	9.65	5.470	...	...	...	BD+14 334, M2
02 07 20.0	50 17 22	8.76	5.760	...	...	...	BD+49 555
02 10 54.1	43 08 30	10.1	0.810	0.490	5.930	1.300	GSC 02842-01596
02 22 44.6	14 49 40	9.67	0.800	0.300	5.180	1.100	BD+14 386, K2
02 24 14.1	40 52 53	10.21	0.750	0.320	6.600	1.070	GSC 02835-00621
02 30 13.0	41 14 53	9.75	0.740	0.500	6.090	1.240	BD+40 535, K2
02 35 37.7	40 35 33	9.89	0.790	0.490	6.090	1.280	GSC 02836-00893
02 47 30.4	-21 58 36	10.0	0.765	0.324	5.707	1.089	BD-22 481 K5
03 01 25.9	-05 02 29	9.3	0.756	0.368	5.671	1.124	BD-05 562
03 02 13.6	-23 48 37	9.9	0.703	0.333	6.129	1.036	CD-24 1384 K5
03 04 31.5	-24 25 17	9.6	0.754	0.311	5.595	1.065	CD-24 1405 K5
03 07 21.6	35 40 41	9.15	6.140	...	5.120	1.020	HD 278277, K2
03 26 25.3	37 48 11	9.75	0.720	0.330	5.570	1.050	HD 275416, M0
03 37 24.0	41 51 22	10.6	0.620	0.430	7.100	1.050	GSC 02870-10994
03 44 17.2	19 05 27	...	0.780	0.640	6.180	1.420	GSC 01256-00228
03 45 29.8	-11 17 41	9.8	0.784	0.307	6.520	1.091	BD-11 729
03 59 12.7	15 48 52	8.90	5.550	...	4.450	1.100	HD 285324, M0
04 02 12.8	21 37 20	10.6	0.730	0.310	6.210	1.040	GSC 01262-00909
04 05 28.1	18 33 22	...	0.770	0.320	5.740	1.090	HD 285353, K5
04 10 46.0	-09 15 58	9.9	0.800	0.329	6.721	1.129	BD-09 838
04 15 51.9	-66 12 32	10.3	0.774	0.347	6.285	1.121	CPD-66 267B
04 20 16.6	18 09 43	9.9	...	...	4.390	...	HD 285651, M0
04 29 46.0	-14 47 07	9.3	0.786	0.309	7.195	1.095	PPM 710748
04 31 25.3	-25 31 02	10.8	0.726	0.346	6.540	1.072	GSC 06467-02288
05 14 26.8	-30 20 57	10.4	0.776	0.318	5.591	1.094	CD-30 2265
05 22 55.8	-17 06 30	9.4	0.796	0.328	6.916	1.124	PPM 711510
05 40 26.7	-16 02 46	10.3	0.744	0.330	6.115	1.074	GSC 05917-01476
05 40 31.3	-17 17 54	10.1	0.748	0.309	6.372	1.057	BD-17 1208
05 53 00.9	-77 18 44	10.5	0.766	0.313	6.212	1.079	PPM 784266
05 55 00.7	-23 35 35	9.7	0.827	0.367	6.167	1.194	CD-23 3232
06 00 25.0	-28 02 28	10.2	0.751	0.303	6.477	1.054	CD-28 2635
06 09 24.8	77 33 27	9.5	0.703	0.453	6.018	1.156	BD+77 225 K0
06 23 31.1	-17 11 13	10.5	0.783	0.317	6.082	1.100	BD-17 141
06 38 29.4	-24 05 09	9.8	0.768	0.301	5.152	1.069	CD-23 4148
06 40 31.0	52 09 16	10.3	0.682	0.352	6.936	1.034	GSC 03389-00656
06 42 35.9	34 55 58	9.5	0.597	0.438	6.406	1.035	GSC 02443-00593
06 49 05.6	34 24 53	9.4	5.88:	...	4.802	1.078	HD 264056 M2
07 00 36.5	26 08 18	10.5	0.844	0.511	6.189	1.355	GSC 01899-00620
07 01 13.8	-37 36 05	10.6	0.753	0.312	6.488	1.065	GSC 07629-00161
07 04 25.6	-41 25 47	10.5	0.794	0.319	6.175	1.113	GSC 07367-00412
07 14 33.4	69 32 22	10.0	0.730	0.362	6.241	1.092	BD+69 412 B5!
07 23 57.1	16 48 29	10.3	0.776	0.310	6.190	1.086	GSC 01347-00105

Table 16—Continued

α (2000)	δ	V	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
07 24 02.7	30 15 45	9.9	0.696	0.350	6.474	1.046	GSC 02452-01602
07 39 19.9	12 36 07	10.7	0.799	0.321	5.813	1.120	GSC 00773-00602
07 50 24.7	22 09 36	10.5	0.794	0.301	5.808	1.095	BD+22 1785
07 52 56.4	38 09 53	10.2	0.800	0.302	5.334	1.102	GSC 02960-00950
08 18 23.2	00 22 52	10.0	0.773	0.310	5.454	1.083	GSC 04848-01624
08 19 18.1	12 04 10	10.1	0.661	0.369	6.469	1.030	GSC 00803-00179
08 49 04.4	-18 34 06	10.1	0.749	0.323	5.936	1.072	BD-18 2490
08 50 22.6	-16 41 39	10.2	0.795	0.330	6.015	1.125	BD-16 2604
08 59 55.5	15 30 16	9.4	0.720	0.320	5.263	1.040	HD 76958
09 23 09.2	-10 03 14	9.4	0.791	0.336	5.868	1.127	PPM 716038
09 47 40.7	-04 55 46	10.1	0.786	0.303	6.096	1.089	BD-04 2721
10 02 37.5	-11 46 14	10.5	0.725	0.304	6.299	1.029	BD-11 2780
10 27 09.8	23 13 47	9.1	5.77:	...	4.732	1.035	BD+23 2226 M0
11 16 30.3	-20 27 33	9.8	0.688	0.317	5.875	1.005	BD-19 3217
11 25 07.5	-29 48 01	10.5	0.740	0.307	6.471	1.047	CD-29 9084
11 30 09.6	17 03 36	9.3	5.64:	...	4.521	1.123	BD+17 2364 M2
11 46 29.2	-27 12 05	9.9	0.789	0.340	5.020	1.129	CD-26 8766
11 58 53.1	-8 24 12	9.0	0.769	0.324	5.253	1.093	BD-07 3328
12 15 49.7	28 47 55	9.4	5.56:	...	4.437	1.121	BD+29 2271
12 26 32.0	65 10 58	10.6	0.769	0.325	6.540	1.094	GSC 04161-00886
13 00 02.5	47 26 32	9.4	0.579	0.557	6.112	1.136	CD CVn K0 III
13 18 18.1	-29 53 25	10.0	0.771	0.302	5.771	1.073	CD-29 10249
13 28 01.0	-07 01 37	9.4	5.79:	...	...	...	HD 117081 K5
13 41 01.8	56 34 52	9.6	0.726	0.450	5.911	1.176	HD 238271 K5
13 47 40.5	-29 58 30	10.0	0.785	0.302	6.006	1.087	CD-29 10597
14 28 46.3	63 19 26	9.5	0.600	0.412	6.323	1.012	BD+63 1135 K2
14 44 37.9	-23 28 23	9.9	0.774	0.310	5.363	1.084	CPD-22 5830
14 44 52.6	-27 12 03	10.7	0.772	0.330	6.710	1.102	GSC 06754-00323
14 57 59.8	-22 51 30	10.9	0.779	0.332	6.516	1.111	GSC 06748-00155
15 11 48.5	-30 50 23	10.4	0.738	0.320	7.063	1.058	CD-30 12022
15 14 23.0	-35 50 47	10.9	0.787	0.310	6.760	1.097	GSC 07324-00450
15 14 43.0	-19 24 25	9.9	0.741	0.302	5.793	1.043	BD-18 4010
15 21 53.1	-13 51 08	10.4	0.798	0.313	5.941	1.111	BD-13 4138
15 27 37.6	-15 57 35	10.3	0.789	0.309	5.158	1.098	BD-15 4113
15 31 01.8	-24 47 16	10.5	0.719	0.313	6.836	1.032	CD-24 12131
15 37 14.5	-23 59 11	9.7	0.797	0.338	6.082	1.135	CD-23 12419
15 45 01.7	-23 23 51	10.4	0.748	0.304	6.099	1.052	BD-22 4016 K5
15 47 34.1	-20 14 34	10.2	0.787	0.311	6.911	1.098	PPM 732355
15 49 19.0	-16 01 22	10.0	0.772	0.309	5.685	1.081	BD-15 4196
16 51 51.0	47 31 14	9.24	...	...	4.700	...	BD+47 2395, M0
17 21 00.1	41 21 48	8.22	...	...	4.360	...	BD+41 2820, K2
17 22 17.9	36 29 08	9.3	5.85:	...	4.792	1.061	BD+36 2860 M2
17 24 21.5	65 29 15	9.6	0.673	0.566	6.157	1.239	BD+65 1182 K2
17 25 20.7	65 06 43	9.9	0.648	0.449	6.429	1.097	GSC 04206-00695
17 26 19.2	60 17 48	9.9	0.701	0.672	5.788	1.373	BD+60 1757 K2
17 31 07.0	49 43 27	9.16	...	...	4.260	...	BD+49 2652
17 36 09.2	52 28 59	8.41	...	...	4.200	...	HD 160270
17 39 26.8	14 19 14	10.0	0.785	0.306	5.224	1.091	GSC 01005-01782
17 57 10.6	26 05 20	11.0	0.777	0.311	6.981	1.088	GSC 02094-00083

Table 16—Continued

α (2000)	δ	V	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
18 04 45.7	18 30 49	10.2	0.759	0.319	6.105	1.078	GSC 01558-00135
18 05 52.9	20 54 01	10.7	0.744	0.302	6.609	1.046	GSC 01566-00162
18 06 09.7	76 38 07	11.2	5.61:	...	4.520	1.086	BD+76 679 K5
18 07 49.9	35 12 29	10.5	0.759	0.349	6.789	1.108	GSC 02629-01650
18 09 42.5	51 14 47	9.1	5.52:	...	4.441	1.086	HD 234573 M0
18 15 48.0	32 23 39	10.1	0.783	0.300	5.540	1.083	GSC 02626-00119
18 16 08.3	21 39 32	8.0	0.777	0.333	5.160	1.110	HD 341857 M0
18 16 21.2	20 28 17	9.1	0.692	0.628	5.135	1.320	HD 348183 K7
18 16 34.2	23 53 27	10.2	0.759	0.301	5.344	1.060	HD 341795 K5
18 19 42.9	27 57 17	9.2	5.57:	...	4.481	1.094	HD 335891 M0
18 32 47.7	76 44 45	10.6	0.779	0.320	5.605	1.099	GSC 04570-00463
19 48 12.5	-31 39 43	9.8	0.765	0.306	5.797	1.071	HD 186874 K1/K2III
19 59 17.0	-27 32 30	9.7	0.792	0.352	6.054	1.144	PPM 735797
20 06 51.7	-01 11 25	10.4	0.753	0.318	6.650	1.071	PPM 708186
20 12 35.0	-02 37 13	9.6	0.776	0.318	5.754	1.094	BD-03 4816
20 14 04.8	-04 03 55	10.4	0.798	0.319	5.634	1.117	BD-04 5062
20 15 14.9	-15 36 26	...	0.819	0.481	6.820	1.300	GSC 06315-00584 var.
20 16 23.7	-04 54 28	10.2	0.787	0.309	5.449	1.096	BD-05 5200
20 18 47.3	-17 27 42	10.4	0.784	0.309	5.833	1.093	BD-17 5932
20 19 52.1	-19 35 58	10.7	0.779	0.365	6.551	1.144	GSC 06336-01895
20 27 31.9	00 22 04	10.9	0.781	0.368	5.515	1.149	GSC 05163-00396
20 28 53.9	-04 05 08	9.5	0.798	0.313	5.540	1.111	BD-04 5150
20 33 16.7	-01 09 34	10.3	0.755	0.310	6.114	1.065	GSC 05176-00531
21 01 55.9	-30 17 07	9.2	0.723	0.300	5.135	1.023	HD 200054 M0 III
21 03 41.5	17 12 54	9.09	0.780	0.360	4.970	1.140	HD 200564, K5
21 06 47.5	17 51 34	11.0	0.750	0.310	6.640	1.060	GSC 01653-01535
21 13 24.2	-35 20 32	10.2	0.646	0.380	6.374	1.026	GSC 07480-00854
21 35 26.7	15 45 03	9.10	0.630	0.520	5.760	1.150	BD+15 4453, K0
21 37 39.3	-23 58 51	10.1	0.820	0.363	5.981	1.183	CD-24 16749
21 42 06.9	-35 20 04	10.3	0.727	0.300	6.735	1.027	CD-35 14918
21 51 51.6	-10 29 35	10.8	0.765	0.316	6.755	1.081	BD-11 5686
21 59 11.1	-29 14 56	10.4	0.739	0.307	6.754	1.046	GSC 06960-01221
22 01 16.7	-01 01 49	10.2	0.800	0.310	5.609	1.110	GSC 05224-01090
22 12 51.4	-47 22 55	11.5	0.780	0.339	6.258	1.119	CD-47 14098
22 17 24.7	-26 20 42	10.8	0.779	0.308	6.385	1.087	GSC 06958-01172
22 28 55.5	-21 36 19	9.8	0.720	0.306	5.912	1.026	HD 213032
22 31 20.1	19 26 16	10.5	0.690	0.340	6.270	1.030	GSC 01703-01220
22 32 00.4	24 52 05	10.1	0.790	0.300	5.440	1.090	GSC 02223-01112
22 56 00.7	41 37 03	9.46	5.870	...	4.790	1.080	BD+40 4943, M0
23 44 37.3	47 24 27	10.36	0.860	0.730	5.150	1.590	GSC 03642-00820

Table 17. Carbon stars

α (2000)	δ	V	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
01 58 51.1	40 37 20	12.1	1.180	0.740	6.630	1.920	CGCS 298
02 13 29.9	46 20 20	13.5	0.860	0.480	6.570	1.340	CGCS 322
03 08 48.7	27 54 59	...	1.030	0.530	6.190	1.560	CGCS 444
03 12 46.9	38 47 50	...	0.970	0.580	6.070	1.550	V458 Per
03 27 31.3	39 04 50	...	1.110	0.530	7.020	1.640	FBS 0324+389
06 14 35.9	-15 01 09	13.0	1.034	0.565	6.382	1.599	CGCS 1210
06 15 27.2	47 54 25	...	1.232	0.838	5.883	2.070	CGCS 1194
07 00 01.7	35 05 56	...	1.126	0.723	6.628	1.849	FBS 0656+351
07 06 07.7	40 12 01	...	1.076	0.523	7.088	1.599	FBS 0702+402
07 54 13.2	72 13 39	13.8	1.050	0.520	5.985	1.570	CGCS 1876
11 45 01.4	64 47 37	...	0.777	0.489	6.034	1.266	GSC 04156-00392
12 24 32.7	-26 14 08	13.6	0.815	0.369	7.219	1.184	KV Hya
20 03 34.2	-25 27 15	12.0	0.783	0.319	7.261	1.102	CD-25 14520

Table 18. Miras and Long-period Variables

α (2000)	δ	V	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
00 09 36.8	37 47 31	12.4	1.010	0.530	5.430	1.540	V414 And
00 38 52.9	45 33 49	...	0.780	0.490	7.710	1.270	V403 And
00 59 46.4	27 56 44	11.4	0.770	0.430	5.980	1.200	W Psc, M2e
01 22 52.9	25 23 06	...	...	...	4.580	1.400	TZ Psc
01 58 06.1	38 39 18	...	0.800	0.360	6.830	1.160	OY And
02 59 56.2	42 37 15	...	0.850	0.630	5.720	1.480	IV Per
03 11 40.7	41 38 46	...	0.840	0.450	5.210	1.290	GG Per
03 18 27.8	42 49 18	...	0.820	0.360	6.230	1.180	WW Per
04 17 41.3	-60 35 42	11.5	0.793	0.474	5.621	1.267	ST Ret
06 14 59.1	46 47 47	11.5	0.873	0.467	4.932	1.340	ST Aur
06 26 59.1	45 47 35	11.5	0.793	0.450	5.505	1.243	BW AUr
06 30 18.3	-60 13 40	10.0	0.800	0.438	5.952	1.238	RU Pic
06 58 33.8	31 38 25	10.0	0.834	0.445	6.295	1.279	FW Gem, M0
07 17 59.7	23 55 19	10.0	0.805	0.444	5.324	1.249	RV Gem
07 56 56.0	31 10 02	11.0	0.784	0.476	5.922	1.260	AO Gem
07 59 14.4	20 38 27	11.0	0.770	0.446	5.019	1.216	BP Gem, M8e
08 03 59.7	73 24 30	10.5	0.801	0.403	5.739	1.204	SW Cam, M5e
08 11 03.1	23 08 54	9.0	5.49:	...	...	...	RR Cnc
08 13 30.6	13 48 05	11.0	0.770	0.327	6.511	1.097	SU Cnc
08 21 53.0	-10 35 50	10.4	0.763	0.502	6.274	1.265	GG Hya
08 38 36.8	-16 17 30	11.0	0.715	0.444	6.739	1.159	FR Hya
09 09 28.6	-22 13 05	13.3	1.088	0.682	4.980	1.770	CC Pyx, M7
09 11 14.5	-09 22 05	11.0	0.695	0.516	5.605	1.211	VV HYa
12 05 14.7	12 21 37	8.4	0.786	0.435	5.357	1.221	SU Vir, M3.5e
12 37 45.8	65 33 21	8.4	0.675	0.335	5.825	1.010	RV Dra, M1
13 59 06.7	-25 51 43	10.0	0.780	0.490	7.007	1.270	FQ HYa
14 38 31.9	-29 43 28	9.8	0.784	0.421	5.372	1.205	FU Hya
14 44 36.9	-19 32 28	10.0	0.804	0.358	5.681	1.162	TW Lib
15 10 44.3	-20 01 08	10.2	0.827	0.493	5.880	1.320	T Lib, M4
15 36 12.7	-21 09 03	10.0	0.783	0.372	6.398	1.155	X Lib
15 37 47.9	-16 09 57	10.3	0.734	0.522	7.199	1.256	W Lib
15 48 38.5	-16 11 41	14.0	0.892	0.745	6.065	1.637	DM Lib
15 59 10.7	-14 10 55	11.0	0.740	0.560	7.650	1.300	UV Lib
16 29 07.8	34 13 46	11.5	0.799	0.312	7.840	1.111	HT Her, ROTSEI
16 59 28.2	12 19 46	10.0	0.810	0.497	4.862	1.307	V440 Oph
17 17 34.2	16 35 26	13.0	0.827	0.470	5.522	1.297	V621 Her
17 41 11.1	12 25 39	12.0	0.888	0.476	6.025	1.364	V1068 Oph
17 52 24.0	34 11 11	12.0	0.760	0.490	6.166	1.250	V1015 Her
17 56 48.2	25 54 21	12.0	0.797	0.327	6.776	1.124	ER Her
18 16 25.0	46 27 51	11.0	0.739	0.383	6.666	1.122	HI Lyr
18 18 20.1	31 42 00	10.0	0.839	0.514	5.113	1.353	AO Lyr
18 19 01.5	22 06 49	14.0	0.862	0.444	7.590	1.306	V577 Her
18 25 59.7	31 29 49	...	0.835	0.419	7.339	1.254	ROTSEI
18 28 56.3	32 14 49	10.0	0.890	0.542	5.692	1.432	IX Lyr
18 30 13.3	37 29 39	10.0	0.826	0.375	6.916	1.201	KL Lyr
18 35 10.0	42 33 34	13.5	5.90:	...	4.577	1.328	ROTSEI
18 35 36.3	39 29 42	...	0.838	0.442	6.256	1.280	ROTSEI
18 42 41.4	35 35 08	12.6	0.789	0.470	7.003	1.259	AX Lyr
18 47 17.8	53 56 47	12.0	0.802	0.464	6.501	1.266	BZ Dra

Table 18—Continued

α (2000)	δ	V	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
18 56 00.5	39 29 16	14.0	1.043	0.674	4.828	1.717	V356 Lyr
18 57 58.0	43 08 05	13.0	0.798	0.467	4.591	1.265	V357 Lyr
19 18 59.4	−23 54 31	12.7	0.737	0.547	6.931	1.284	V1264 Sgr
19 20 12.7	−40 52 37	12.2	0.837	0.534	5.601	1.371	V1158 Sgr
19 22 12.5	−21 41 03	13.2	0.840	0.457	6.546	1.297	V2141 Sgr
19 23 21.1	−23 53 33	13.2	0.829	0.476	6.395	1.305	V1269 Sgr
19 33 55.5	−22 49 44	11.0	0.839	0.484	5.827	1.323	V1315 Sgr, M3e
19 37 52.2	−14 08 08	12.0	0.831	0.553	6.287	1.384	EZ Sgr
19 38 16.3	−32 16 48	11.0	0.822	0.619	6.876	1.441	DP Sgr
19 42 17.3	−37 35 25	12.5	0.859	0.455	6.919	1.314	V2165 Sgr
19 45 42.3	−26 35 26	11.5	0.838	0.502	5.906	1.340	V1167 Sgr
20 05 11.8	−02 39 59	13.0	0.969	0.523	4.841	1.492	V901 Aql
20 07 37.3	−02 27 06	13.0	0.907	0.542	7.080	1.449	V510 Aql
20 08 38.6	−04 25 20	12.5	0.931	0.519	4.693	1.450	V580 Aql
20 10 00.3	−01 41 05	12.0	0.873	0.588	6.539	1.461	V583 Aql.
20 14 29.0	−21 58 44	11.0	0.825	0.479	5.272	1.304	W Cap, M5
20 14 39.9	−01 10 36	12.0	0.895	0.521	5.670	1.416	V519 Aql
20 23 47.5	−03 51 40	14.5	0.780	0.411	5.594	1.191	SON 4473, M3e
20 32 28.2	−05 17 12	12.0	0.724	0.431	6.596	1.155	V837 Aql
20 40 37.0	13 27 00	12.0	0.787	0.514	5.865	1.301	SS Del
21 06 42.6	−01 20 05	9.0	0.782	0.564	5.726	1.346	TX Aqr
21 09 46.7	13 44 17	...	0.820	0.410	4.860	1.230	HK Peg
21 26 46.4	16 34 53	9.0	0.710	0.330	5.120	1.040	TV Peg
21 49 19.0	20 37 46	...	0.820	0.560	4.640	1.380	EL Peg
21 49 31.4	22 41 45	12.0	0.850	0.430	6.740	1.280	CX Peg
22 03 33.9	14 00 31	...	0.790	0.400	6.040	1.190	DG Peg
22 05 59.5	35 30 05	...	0.820	0.530	5.900	1.350	XX Peg
22 12 50.9	−21 09 51	12.0	0.837	0.447	6.484	1.284	AQ Aqr
22 43 17.4	43 40 28	...	0.770	0.630	6.550	1.400	ST Lac
22 53 30.7	−32 55 40	11.0	0.771	0.527	6.726	1.298	SS PsA
23 17 59.6	46 45 12	...	0.860	0.540	5.470	1.400	AO And

Table 19. Semi-regular Variables

α (2000)	δ	V	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
00 45 01.1	48 41 02	10.7	1.250	0.720	5.300	1.970	V864 Cas
00 46 24.8	47 41 33	...	1.280	0.970	4.990	2.250	GSC 03266-01510
02 34 31.4	40 04 06	...	0.830	0.450	7.330	1.280	GU And
03 21 10.4	-17 13 56	10.0	0.693	0.371	6.960	1.064	CO Eri
08 22 01.4	13 37 05	10.8	0.799	0.362	5.833	1.161	GT Cnc
08 27 40.4	19 15 43	11.4	0.730	0.369	6.718	1.099	GV Cnc
08 28 08.0	38 20 22	11.1	0.780	0.353	6.962	1.133	RX Lyn
09 06 39.0	-19 18 44	12.7	1.238	0.843	6.009	2.081	V379 Hya
15 06 42.1	-23 25 38	12.0	0.761	0.520	6.082	1.281	WZ Lib,
16 23 13.1	44 08 28	10.5	0.786	0.318	7.623	1.104	AY Her
19 22 42.9	-32 08 03	9.6	0.733	0.322	5.987	1.055	AM Sgr
19 32 44.7	-32 14 14	12.5	0.839	0.448	5.047	1.287	V347 Sgr
20 51 00.6	14 31 15	12.0	0.800	0.335	7.616	1.135	DZ Del
21 15 02.8	-09 46 37	...	0.876	0.491	5.907	1.367	BY Aqr
23 21 05.5	45 24 30	...	0.750	0.320	7.520	1.070	V338 And
23 25 41.7	45 42 04	...	1.480	0.960	6.280	2.440	V339 And

Table 20. Other Late-type Stars

α (2000)	δ	V	$J - H$	$H - K_S$	K_S	$J - K_S$	ID
04 27 42.7	-73 11 08	...	0.795	0.316	7.742	1.111	WOH S 11, M
08 53 26.7	-13 31 16	...	0.802	0.374	6.566	1.176	V357 Hya, M-type variable
09 31 01.7	-23 09 00	11.0	0.686	0.319	7.526	1.005	AT Hya Irr
09 56 32.9	-10 01 17	15.0	0.846	0.660	5.755	1.506	BR B0954-0947, M8 III
12 44 42.0	36 45 50	9.8	0.797	0.420	6.198	1.217	TX CVn symbiotic
15 01 12.8	-25 39 24	11.6	0.917	0.493	7.126	1.410	BV 1671, variable
15 12 15.2	-22 59 07	12.0	0.856	0.438	5.690	1.294	StM 218, M6
17 25 13.0	64 26 12	...	0.867	0.471	6.603	1.338	FBS 1724+644, M
17 25 42.3	19 33 26	13.0	5.92:	...	4.630	1.299	V400 Her, pulsating variable
18 21 09.5	46 08 57	...	0.680	0.475	6.137	1.155	Rotsei M-type variable
19 36 20.0	-18 29 18	12.2	0.821	0.657	6.524	1.478	V3882 Sgr, variable
19 44 48.9	-15 11 20	13.0	0.901	0.577	6.737	1.478	V362 Sgr, variable
19 51 51.7	-05 48 16	12.7	0.965	0.652	5.328	1.617	EF Aql, variable
19 52 52.4	-38 10 03	12.0	0.777	0.341	6.593	1.118	red giant
19 56 56.8	-11 48 05	13.0	0.936	0.507	5.788	1.443	FI Sgr, variable
20 01 57.7	-03 42 39	...	0.975	0.552	6.599	1.527	Son 8258, variable
20 10 21.2	00 13 26	12.0	0.913	0.740	7.006	1.653	EI Aql, variable
20 24 32.9	-29 44 02	...	0.733	0.329	7.391	1.062	DENIS-P J202432.9-294402, BV 1735

Table 21. $J < 9$ Sources Without a Cataloged Counterpart

α (2000)	δ	$J - H$	$H - K_S$	K_S	$J - K_S$	R_{USNO}
00 11 11.0	48 58 43	0.800	0.320	5.750	1.120	10.9
00 12 10.5	46 53 30	0.800	0.310	6.700	1.110	10.8
00 14 40.9	40 34 52	0.720	0.460	6.350	1.180	10.2
00 19 29.8	48 38 27	0.790	0.310	7.800	1.100	11.8
00 21 48.9	44 52 18	0.800	0.420	7.490	1.220	9.1
00 28 26.9	31 24 11	0.800	0.310	6.630	1.110	9.9
00 38 59.3	-19 27 42	0.788	0.355	6.733	1.143	10.9
00 57 50.8	50 50 07	0.900	0.450	7.370	1.350	13.9
01 00 07.8	46 11 36	0.800	0.310	7.790	1.110	12.1
01 03 57.5	20 11 45	0.770	0.300	6.990	1.070	11.7
01 07 03.8	22 16 06	0.770	0.300	7.260	1.070	10.6
01 08 41.9	48 20 03	0.750	0.310	6.980	1.060	10.8
01 10 50.8	29 17 27	0.780	0.330	7.820	1.110	11.6
01 28 21.2	50 14 51	0.790	0.310	6.180	1.100	9.5
02 08 02.2	41 13 08	0.760	0.300	7.630	1.060	11.1
02 13 41.9	43 25 32	0.760	0.300	6.620	1.060	11.6
02 23 47.8	23 03 41	0.790	0.310	7.070	1.100	11.0
02 36 36.8	30 16 20	0.800	0.300	7.040	1.100	10.4
02 53 46.6	29 32 48	0.750	0.300	7.290	1.050	10.7
03 17 22.1	24 07 02	0.640	0.450	6.550	1.090	10.6
04 02 14.1	20 57 40	0.790	0.340	7.060	1.130	11.2
04 06 59.5	20 23 22	0.770	0.300	7.280	1.070	11.2
04 13 56.1	19 23 54	0.780	0.300	7.050	1.080	10.9
04 14 06.6	19 45 41	0.740	0.370	7.250	1.110	11.7
04 21 20.4	-02 50 02	0.805	0.422	6.960	1.227	12.9
04 27 05.9	00 22 53	0.793	0.300	6.602	1.093	9.5
04 37 43.9	-25 20 34	0.726	0.330	7.306	1.056	10.7
05 03 25.2	73 49 21	0.860	0.438	6.498	1.298	10.9
05 19 11.1	-25 36 46	0.768	0.312	6.800	1.080	11.3
05 22 41.0	-12 08 29	0.796	0.322	7.636	1.118	10.2
05 37 45.4	-23 54 39	0.856	0.428	7.493	1.284	13.3
05 40 50.4	-21 57 04	0.774	0.311	6.793	1.085	9.9
05 45 18.1	-19 57 07	0.789	0.363	6.058	1.152	9.8
05 47 28.1	-21 47 23	1.014	0.550	7.111	1.564	11.3
06 10 32.1	-24 49 27	0.729	0.339	6.935	1.068	10.4
06 13 45.0	52 25 40	0.868	0.508	7.295	1.376	13.8
06 15 36.6	-18 46 52	0.766	0.331	7.260	1.097	10.9
06 18 27.5	-35 25 27	0.798	0.310	6.202	1.108	9.9
06 20 52.1	-16 45 41	0.896	0.460	6.661	1.356	14.7
06 21 44.5	-18 54 21	0.798	0.316	7.615	1.114	11.1
06 28 12.7	-21 24 15	0.807	0.406	6.966	1.213	11.7
06 29 53.0	-21 22 59	0.788	0.329	6.403	1.117	13.0
06 30 46.6	-76 43 09	0.635	0.366	7.915	1.001	...
06 34 20.3	-39 35 18	0.789	0.432	6.941	1.221	14.2
06 35 01.3	-21 40 10	0.795	0.342	6.374	1.137	10.0
06 37 04.8	-40 56 35	0.785	0.314	7.683	1.099	11.7
06 39 59.2	-34 24 16	0.764	0.313	7.511	1.077	11.1
06 40 23.4	-41 28 04	0.730	0.333	7.103	1.063	10.5
06 41 10.1	-35 59 43	0.745	0.326	6.434	1.071	10.3

Table 21—Continued

α (2000)	δ	$J - H$	$H - K_S$	K_S	$J - K_S$	R_{USNO}
06 41 40.3	−28 21 02	0.834	0.452	6.761	1.286	12.6
06 41 57.5	−40 29 44	0.798	0.329	6.393	1.127	10.8
06 45 16.7	−30 08 35	0.735	0.315	6.646	1.050	11.6
06 50 54.8	−37 29 22	0.950	0.511	7.118	1.461	15.0
06 51 13.3	−37 56 49	0.817	0.365	6.580	1.182	10.9
06 51 55.8	−41 04 40	0.780	0.395	7.665	1.175	11.5
06 52 22.8	45 20 45	0.874	0.661	6.147	1.535	11.3
06 53 50.7	−37 41 56	0.828	0.370	6.693	1.198	11.2
06 55 08.7	−37 57 18	0.761	0.301	6.925	1.062	10.1
06 57 52.0	66 21 11	0.861	0.556	6.103	1.417	11.4
06 58 11.8	26 35 35	1.187	0.672	6.713	1.859	11.4
07 01 18.8	38 30 01	0.838	0.432	7.362	1.270	12.8
07 01 32.2	−38 14 21	0.877	0.468	6.847	1.345	12.2
07 03 35.6	−40 47 48	0.846	0.454	7.059	1.300	12.6
07 04 17.0	24 57 47	0.777	0.308	6.255	1.085	10.4
07 06 53.0	40 47 29	0.793	0.317	5.454	1.110	10.3
07 10 48.3	30 55 46	1.022	0.546	6.588	1.568	11.0
07 10 57.4	47 58 18	1.056	0.530	7.354	1.586	12.3
07 17 09.5	27 48 04	0.782	0.304	7.261	1.086	11.6
07 21 40.4	19 43 50	0.900	0.464	6.497	1.364	11.4
07 24 16.5	14 14 56	0.791	0.370	6.934	1.161	10.6
07 27 47.5	18 14 37	0.838	0.373	5.616	1.211	12.0
07 30 31.4	22 36 55	0.834	0.433	5.699	1.267	12.4
07 31 35.6	41 05 47	0.760	0.301	6.303	1.061	9.9
07 36 34.6	15 05 00	0.787	0.319	7.015	1.106	10.8
07 56 36.3	18 52 16	0.791	0.367	7.181	1.158	11.5
08 01 01.6	00 05 22	0.797	0.324	7.251	1.121	11.4
08 01 48.5	50 42 43	0.784	0.300	6.565	1.084	10.9
08 05 47.9	52 46 28	0.838	0.375	7.142	1.213	12.1
08 09 24.4	−05 53 29	0.787	0.313	5.444	1.100	9.0
08 10 17.8	−04 57 16	0.731	0.367	7.770	1.098	11.9
08 13 34.3	−05 13 21	1.230	0.714	6.910	1.944	11.8
08 22 44.9	16 31 39	0.763	0.327	6.716	1.090	11.5
08 26 30.0	−04 05 32	0.805	0.366	5.833	1.171	10.7
08 28 06.9	−11 25 56	0.736	0.311	7.567	1.047	11.5
08 29 15.1	18 23 07	1.033	0.604	7.090	1.637	11.5
08 30 06.0	−12 15 40	0.819	0.365	7.495	1.184	11.7
08 46 25.3	−18 21 44	0.729	0.354	6.125	1.083	9.5
08 51 16.6	−07 27 32	0.811	0.367	6.578	1.178	12.0
08 55 40.2	−17 26 42	0.794	0.308	6.985	1.102	10.4
09 03 25.2	−21 53 12	0.774	0.404	6.239	1.178	9.9
09 07 31.8	−07 42 25	0.805	0.369	5.114	1.174	9.7
09 09 35.6	−10 31 00	0.750	0.313	7.613	1.063	11.1
09 28 22.3	−19 49 04	0.742	0.312	7.113	1.054	10.8
09 30 46.5	−19 32 41	0.798	0.341	6.789	1.139	11.4
09 34 56.4	−23 47 55	0.779	0.300	6.462	1.079	9.7
09 42 23.5	−09 47 50	0.799	0.320	6.354	1.119	10.4
09 42 45.1	−20 49 02	0.742	0.305	7.478	1.047	10.9
09 54 35.4	−19 26 05	0.781	0.338	6.284	1.119	11.1

Table 21—Continued

α (2000)	δ	$J - H$	$H - K_S$	K_S	$J - K_S$	R_{USNO}
10 04 32.4	−24 35 53	0.779	0.356	6.413	1.135	11.1
10 10 01.5	−02 37 43	0.824	0.465	7.008	1.289	11.4
10 42 05.0	−08 20 28	0.784	0.312	6.971	1.096	10.6
11 05 44.9	−21 38 23	0.699	0.314	6.672	1.013	10.58
11 28 41.0	−17 10 05	0.801	0.366	6.831	1.167	12.3
11 46 47.0	23 51 20	0.760	0.301	5.792	1.061	10.7
11 58 16.9	−25 37 53	0.774	0.497	7.098	1.271	11.4
12 00 28.5	64 09 39	0.798	0.353	6.129	1.151	11.8
12 08 50.4	−28 10 14	0.766	0.373	7.423	1.139	11.1
12 13 12.0	−26 23 47	0.703	0.300	6.907	1.003	10.7
12 21 01.1	−19 52 38	0.794	0.307	6.872	1.101	10.4
12 24 42.4	−27 30 14	0.781	0.326	6.695	1.107	11.0
13 18 50.8	−17 59 50	0.766	0.333	7.545	1.099	11.5
13 26 21.9	−19 05 23	0.744	0.309	6.914	1.053	15.1
13 29 03.1	−45 38 36	0.793	0.338	6.678	1.131	11.1
14 27 23.3	−35 29 25	0.799	0.329	6.431	1.128	10.1
14 48 00.2	−25 49 10	0.776	0.342	7.869	1.118	12.2
14 48 05.4	−23 10 37	0.792	0.305	6.383	1.097	10.1
14 50 21.1	−35 09 03	0.794	0.312	7.728	1.106	11.5
14 50 44.0	−19 27 04	0.758	0.300	7.080	1.058	10.6
14 51 17.2	−31 31 28	0.800	0.306	7.263	1.106	11.3
14 59 09.5	−27 02 49	0.721	0.301	7.484	1.022	10.8
14 59 19.9	−35 48 06	0.776	0.321	6.207	1.097	10.5
15 02 09.9	59 31 21	0.914	0.476	7.316	1.390	13.7
15 02 37.9	−35 16 43	0.788	0.321	7.881	1.109	14.0
15 02 57.1	−03 49 45	0.796	0.311	7.211	1.107	11.0
15 02 58.2	−35 51 11	0.849	0.515	7.299	1.364	14.1
15 04 40.3	−31 46 50	0.788	0.304	6.455	1.092	10.1
15 04 46.8	−32 17 18	0.782	0.332	6.311	1.114	10.3
15 06 15.1	−30 09 23	0.739	0.344	7.509	1.083	11.1
15 06 59.4	−18 39 51	0.797	0.307	6.470	1.104	9.8
15 12 11.9	−35 48 29	0.695	0.357	7.408	1.052	10.7
15 13 56.0	−30 56 05	0.884	0.445	7.024	1.329	...
15 43 34.5	−19 25 04	0.746	0.415	6.413	1.161	11.0
15 53 03.2	−19 59 22	0.772	0.347	6.777	1.119	10.7
16 04 56.0	−19 01 57	0.760	0.347	7.248	1.107	11.0
16 25 01.3	49 28 16	0.715	0.303	6.868	1.018	10.8
16 55 50.4	31 41 25	0.785	0.317	6.870	1.102	11.0
17 02 54.2	33 42 55	0.786	0.346	6.840	1.132	10.3
17 05 49.2	35 17 04	0.793	0.356	6.611	1.149	11.1
18 00 36.3	13 17 17	0.793	0.316	6.434	1.109	10.3
18 07 10.8	53 42 52	0.775	0.335	7.071	1.110	10.9
18 53 53.0	−47 19 17	0.770	0.333	6.294	1.103	10.4
19 21 29.8	−23 00 23	0.756	0.335	6.202	1.091	9.7
19 21 34.7	−23 06 55	0.796	0.307	6.717	1.103	10.6
19 22 39.6	−32 39 35	0.792	0.306	6.492	1.098	10.0
19 23 14.7	−22 21 54	0.764	0.317	7.061	1.081	11.0
19 23 48.8	−41 56 25	0.810	0.362	7.301	1.172	12.2
19 24 58.3	−33 08 49	0.783	0.312	7.018	1.095	10.7

Table 21—Continued

α (2000)	δ	$J - H$	$H - K_S$	K_S	$J - K_S$	R_{USNO}
19 26 30.6	−19 56 55	0.796	0.325	7.176	1.121	11.0
19 29 00.9	−20 42 32	0.684	0.361	7.447	1.045	10.8
19 29 22.3	−22 02 26	0.793	0.312	6.767	1.105	10.6
19 30 15.5	−23 20 48	0.769	0.494	7.730	1.263	15.9
19 30 23.8	−23 42 45	0.758	0.305	6.948	1.063	10.4
19 31 39.8	−17 57 36	0.742	0.400	7.431	1.142	11.2
19 31 41.9	−20 36 23	0.758	0.348	7.391	1.106	11.6
19 34 18.2	−32 32 58	0.818	0.418	7.094	1.236	11.8
19 37 14.0	−24 48 01	0.784	0.308	7.279	1.092	11.4
19 37 21.8	−17 26 42	0.795	0.360	7.242	1.155	11.4
19 40 29.1	−35 38 08	0.763	0.333	6.954	1.096	10.6
19 40 43.7	−32 36 09	0.862	0.437	7.692	1.299	14.8
19 41 28.4	−20 50 50	0.790	0.361	6.524	1.151	11.4
19 42 44.1	−29 54 36	0.883	0.458	7.315	1.341	12.2
19 43 46.7	−18 52 11	0.789	0.314	6.811	1.103	10.4
19 44 53.1	−20 05 27	0.747	0.350	7.555	1.097	11.2
19 46 45.5	−28 05 51	0.760	0.331	6.938	1.091	10.3
19 48 36.9	−19 58 39	0.725	0.332	6.927	1.057	10.8
19 49 06.4	−20 22 22	0.772	0.333	7.321	1.105	11.0
19 50 33.0	−36 41 06	0.701	0.308	7.017	1.009	11.2
19 51 36.4	−24 04 12	0.772	0.364	6.767	1.136	10.9
19 51 43.3	−24 00 52	0.804	0.355	7.796	1.159	11.9
19 56 40.9	−24 43 40	0.749	0.301	7.127	1.050	10.4
20 08 51.5	−27 02 02	0.760	0.303	7.506	1.063	11.1
20 10 23.7	−19 44 45	0.744	0.334	7.029	1.078	11.0
20 11 09.1	−04 01 00	0.821	0.419	7.432	1.240	12.5
20 12 41.1	−23 41 34	0.773	0.300	7.483	1.073	11.0
20 15 05.3	−05 20 52	0.775	0.321	6.146	1.096	10.9
20 16 21.7	00 34 15	0.705	0.304	6.761	1.009	10.7
20 17 37.9	−27 02 05	0.786	0.321	6.780	1.107	11.1
20 17 49.6	−14 15 33	0.696	0.307	7.770	1.003	11.4
20 20 47.4	00 54 12	0.763	0.304	7.436	1.067	11.3
20 22 18.9	−27 10 40	0.791	0.316	7.528	1.107	11.7
20 25 46.4	−16 31 48	0.826	0.497	6.492	1.323	11.0
20 27 43.1	−39 22 18	0.772	0.304	6.557	1.076	10.4
20 29 04.5	−22 57 23	0.758	0.319	6.563	1.077	9.5
20 31 34.2	−02 17 53	0.776	0.344	6.069	1.120	10.5
20 33 36.9	−04 01 26	0.876	0.440	6.270	1.316	11.8
20 39 49.1	−23 55 31	0.785	0.357	6.317	1.142	10.3
20 44 54.0	−07 43 59	0.836	0.487	7.418	1.323	14.2
20 45 12.5	−04 37 22	0.782	0.305	7.248	1.087	10.9
20 46 14.6	−03 31 50	0.795	0.308	7.365	1.103	11.4
20 50 55.8	−21 00 27	0.761	0.343	6.279	1.104	9.9
21 05 48.8	13 59 01	0.790	0.360	7.410	1.150	11.9
21 16 28.6	20 22 00	0.780	0.310	6.870	1.090	10.5
21 25 44.0	16 02 10	0.740	0.360	7.060	1.100	11.9
21 44 53.3	−04 05 46	0.759	0.343	6.867	1.102	10.2
22 34 35.1	45 04 15	0.850	0.510	7.460	1.360	15.5
22 42 31.5	35 06 43	0.740	0.320	7.820	1.060	12.1

Table 21—Continued

α (2000)	δ	$J - H$	$H - K_S$	K_S	$J - K_S$	R_{USNO}
22 42 41.3	44 41 04	0.830	0.370	5.860	1.200	11.6
23 09 19.6	47 32 18	0.790	0.450	6.800	1.240	10.9
23 34 07.6	46 20 02	0.950	0.520	6.670	1.470	14.0
23 34 20.7	42 23 08	0.770	0.310	7.760	1.080	12.3
23 37 37.5	41 53 37	0.820	0.370	5.800	1.190	11.2
23 50 42.0	23 19 41	0.860	0.480	7.140	1.340	14.2